arXiv:0707.0165v1 [math.GR] 2 Jul 2007

[1]

ALGORITHMIC PROBLEMS IN AMALGAMS OF FINITE GROUPS: CONJUGACY AND INTERSECTION PROPERTIES

L.MARKUS-EPSTEIN

Abstract. Geometric methods proposed by Stallings [46] for treating finitely generated subgroups of free groups were successfully used to solve a wide collection of decision problems for free groups and their subgroups [3, 19, 29, 30, 36, 41, 49].

In the present paper we employ the generalized Stallings' methods, developed by the author in [32], to solve various algorithmic problems concerning finitely generated subgroups of amalgams of finite groups.

1. Introduction

This paper continues the line of [33] and [34]. The primary goal of the sequence of these three papers is to solve effectively (by finding an algorithm) various decision problems concerning finitely generated subgroups of amalgams of finite groups.

Decision (or *algorithmic*) problems is one of the classical subjects of combinatorial group theory originating in the three fundamental decision problems posed by Dehn [9] in 1911: the word problem, the conjugacy problem and the isomorphism problem. As is well known (the reader is referred to [37, 38] for a survey on decision problems for groups), these problems are theoretically undecidable in general. Thus the celebrated Novikov-Boone theorem asserts that the word problem is undecidable (p.88 in [27]). However restrictions to some particular classes of groups may yield surprisingly good results. Remarkable examples include the solvability of the word problem in one-relator groups (Magnus, see II.5.4 in [27]) and in hyperbolic groups (Gromov, see 2.3.B in [15]).

In free groups a big success in this direction is due to the geometrical methods proposed by Stallings [46] in the early 80's. Recall that Stallings showed that every finitely generated subgroup of a free group is canonically represented by a minimal immersion of a bouquet of circles. Using the graph theoretic language, the results of [46] can be restated as follows. A finitely generated subgroup of a free group is canonically represented by a finite labelled graph which can be constructed algorithmically by a so called

---

[1] Supported in part at the Technion by a fellowship of the Israel Council for Higher Education

process of *Stallings' foldings* (*Stallings' folding algorithm*). Moreover, this algorithm is quadratic in the size of the input [19, 30]. See [48] for a faster implementation of this algorithm.

This approach reviled as extremely useful to solve algorithmic problems in free groups. See [3, 29, 30, 36, 41, 49] for examples of the applications of the Stallings' approach in free groups, and [22, 24, 35, 42] for the applications in some other classes of groups. Note that Stallings' ideas were recast in a combinatorial graph theoretic way in the remarkable survey paper of Kapovich and Myasnikov [19], where these methods were applied systematically to study the subgroup structure of free groups.

Our recent results [32] show that Stallings' methods can be completely generalized to the class of amalgams of finite groups. Along the current paper we refer to this generalization of Stallings' folding algorithm as the *generalized Stallings' folding algorithm*. Its description is included in the Appendix. Let us emphasize that the generalized Stallings' algorithm is quadratic in the size of the input, which yields a quadratic time solution of the membership problem in amalgams of finite groups (see [32]).

We employ these generalized Stallings' methods to answer a collection of algorithmic questions concerning finitely generated subgroups of amalgams of finite groups, which extends the results presented in [19]. Our results include polynomial solutions for the following algorithmic problems (which are known to be unsolvable in general [37, 38]) in amalgams of finite groups:

- computing subgroup presentations,
- detecting triviality of a given subgroup,
- the freeness problem,
- the finite index problem,
- the separability problem,
- the conjugacy problem,
- the normality,
- the intersection problem,
- the malnormality problem,
- the power problem,
- reading off Kurosh decomposition for finitely generated subgroups of free products of finite groups.

These results are spread out between three papers: [33, 34] and the current one. In [34] free products of finite groups are considered, and an efficient procedure to read off a Kurosh decomposition is presented.

The splitting between [33] and the current paper was done with the following idea in mind. It turn out that some subgroup properties, such as computing of a subgroup presentation and index, as well as detecting of freeness and normality, can be obtained directly by an analysis of the corresponding subgroup graph. Solutions of others require some additional constructions. Thus, for example, intersection properties can be examined via product graphs, and separability needs constructions of a pushout of graphs.

In [33] algorithmic problems of the first type are presented: the computing of subgroup presentations, the freeness problem and the finite index problem. The separability problem is also included there, because it is closely related with the other problems presented in [33]. The rest of the algorithmic problems are introduced in the current paper.

The paper is organized as follows. The Preliminary Section includes the description of the basic notions used along the present paper. Readers familiar with amalgams, normal words in amalgams and labelled graphs can skip it. The next section presents a summary of the results from [32] which are essential for our algorithmic purposes. It describes the nature and the properties of the subgroup graphs constructed by the generalized Stallings' folding algorithm in [32]. The rest of the sections are titled by the names of various algorithmic problems and present definitions (descriptions) and solutions of the corresponding algorithmic problems. The relevant references to other papers considering similar problems and a rough analysis of the complexity of the presented solutions (algorithms) are provided. In contrast with papers that establish the exploration of the complexity of decision problems as their main goal (for instance, [20, 21, 48]), we do it rapidly (sketchy) viewing in its analysis a way to emphasize the effectiveness of our methods.

Other Methods. There have been a number of papers, where methods, not based on Stallings' foldings, have been presented. One can use these methods to treat finitely generated subgroups of amalgams of finite groups. A topological approach can be found in works of Bogopolskii [4, 5]. For the automata theoretic approach, see papers of Holt and Hurt [17, 18], papers of Cremanns, Kuhn, Madlener and Otto [8, 25], as well as the recent paper of Lohrey and Senizergues [26].

However the methods for treating finitely generated subgroups presented in the above papers were applied to some particular subgroup property. No one of these papers has as its goal a solution of various algorithmic problems, which we consider as our primary aim. Moreover, similarly to the case of free groups (see [19]), our combinatorial approach seems to be the most natural one for this purpose.

2. Acknowledgments

I wish to deeply thank to my PhD advisor Prof. Stuart W. Margolis for introducing me to this subject, for his help and encouragement throughout my work on the thesis. I owe gratitude to Prof. Arye Juhasz for his suggestions and many useful comments during the writing of this paper. I gratefully acknowledge a partial support at the Technion by a fellowship of the Israel Council for Higher Education.

3. Preliminaries

Amalgams. Let $G = G_1 \ast_A G_2$ be a free product of G_1 and G_2 with amalgamation, customary, an *amalgam* of G_1 and G_2. We assume that the (free) factors are given by the finite group presentations

$$G_1 = gp\langle X_1 | R_1 \rangle, \quad G_2 = gp\langle X_2 | R_2 \rangle \ \text{ such that } \ X_1^{\pm} \cap X_2^{\pm} = \emptyset. \tag{1.a}$$

$A = \langle Y \rangle$ is a group such that there exist two monomorphisms

$$\phi_1 : A \to G_1 \text{ and } \phi_2 : A \to G_2. \tag{1.b}$$

Thus G has a finite group presentation

$$G = gp\langle X_1, X_2 | R_1, R_2, \phi_1(a) = \phi_2(a), \ a \in Y \rangle. \tag{1.c}$$

We put $X = X_1 \cup X_2$, $R = R_1 \cup R_2 \cup \{\phi_1(a) = \phi_2(a) \mid a \in Y\}$. Thus $G = gp\langle X | R \rangle$.

As is well known [27, 28, 43], the free factors embed in G. It enables us to identify A with its monomorphic image in each one of the free factors. Sometimes in order to make the context clear we use $\boxed{G_i \cap A}$ [1] to denote the monomorphic image of A in G_i ($i \in \{1, 2\}$).

Elements of $G = gp\langle X | R \rangle$ are equivalence classes of words. However it is customary to blur the distinction between a word u and the equivalence class containing u. We will distinguish between them by using different equality signs: $\boxed{\text{“}\equiv\text{”}}$ for the equality of two words and $\boxed{\text{“}=_G\text{”}}$ to denote the equality of two elements of G, that is the equality of two equivalence classes. Thus in $G = gp\langle x \mid x^4 \rangle$, for example, $x \equiv x$ but $x \not\equiv x^{-3}$, while $x =_G x^{-3}$.

Normal Forms. Let $G = G_1 \ast_A G_2$. A word $g_1 g_2 \cdots g_n \in G$ is *in normal form* (or, simply, it is a *normal word*) if:

1. $g_i \neq_G 1$ lies in one of the factors, G_1 or G_2,
2. g_i and g_{i+1} are in different factors,
3. if $n \neq 1$, then $g_i \notin A$.

---

[1] Boxes are used for emphasizing purposes only.

We call the sequence $(g_1, g_2, \ldots, g_n)$ a *normal decomposition* of the element $g \in G$, where $g =_G g_1 g_2 \cdots g_n$.

Any $g \in G$ has a representative in a normal form (see, for instance, p.187 in [27]). If $g \equiv g_1 g_2 \cdots g_n$ is in normal form and $n > 1$, then the Normal Form Theorem (IV.2.6 in [27]) implies that $g \neq_G 1$. The number n is unique for a given element g of G and it is called the *syllable length* of g. We denote it $l(g)$. We use $|g|$ to denote the length of g as a word in X^*.

Labelled graphs. Below we follow the notation of [11, 46].

A graph Γ consists of two sets $E(\Gamma)$ and $V(\Gamma)$, and two functions $E(\Gamma) \to E(\Gamma)$ and $E(\Gamma) \to V(\Gamma)$: for each $e \in E$ there is an element $\overline{e} \in E(\Gamma)$ and an element $\iota(e) \in V(\Gamma)$, such that $\overline{\overline{e}} = e$ and $\overline{e} \neq e$.

The elements of $E(\Gamma)$ are called *edges*, and an $e \in E(\Gamma)$ is a *direct edge* of Γ, $\overline{e}$ is the *reverse (inverse) edge* of e.

The elements of $V(\Gamma)$ are called *vertices*, $\iota(e)$ is the *initial vertex* of e, and $\tau(e) = \iota(\overline{e})$ is the *terminal vertex* of e. We call them the *endpoints* of the edge e.

A *path of length* n is a sequence of n edges $p = e_1 \cdots e_n$ such that $v_i = \tau(e_i) = \iota(e_{i+1})$ $(1 \leq i < n)$. We call p a *path from* $v_0 = \iota(e_1)$ *to* $v_n = \tau(e_n)$. The *inverse* of the path p is $\overline{p} = \overline{e_n} \cdots \overline{e_1}$. A path of length 0 is the *empty path*.

We say that the graph Γ is *connected* if $V(\Gamma) \neq \emptyset$ and any two vertices are joined by a path. The path p is *closed* if $\iota(p) = \tau(p)$, and it is *freely reduced* if $e_{i+1} \neq \overline{e_i}$ $(1 \leq i < n)$. Γ is a *tree* if it is a connected graph and every closed freely reduced path in Γ is empty.

A *subgraph* of Γ is a graph C such that $V(C) \subseteq V(\Gamma)$ and $E(C) \subseteq E(\Gamma)$. In this case, by abuse of language, we write $C \subseteq \Gamma$. Similarly, whenever we write $\Gamma_1 \cup \Gamma_2$ or $\Gamma_1 \cap \Gamma_2$, we always mean that the set operations are, in fact, applied to the vertex sets and the edge sets of the corresponding graphs.

A *labelling* of Γ by the set $X^{\pm}$ is a function

$$lab: \ E(\Gamma) \to X^{\pm}$$

such that for each $e \in E(\Gamma)$, $lab(\overline{e}) \equiv (lab(e))^{-1}$.

The last equality enables one, when representing the labelled graph Γ as a directed diagram, to represent only X-labelled edges, because X^{-1}-labelled edges can be deduced immediately from them.

A graph with a labelling function is called a *labelled (with $X^{\pm}$) graph*. The only graphs considered in the present paper are labelled graphs.

A labelled graph is called *well-labelled* if

$$\iota(e_1) = \iota(e_2), \ lab(e_1) \equiv lab(e_2) \ \Rightarrow \ e_1 = e_2,$$

for each pair of edges $e_1, e_2 \in E(\Gamma)$. See Figure 1.



FIGURE 1. The graph Γ_1 is labelled with $\{a, b, c\}^{\pm}$, but it is not well-labelled. The graphs Γ_2 and Γ_3 are well-labelled with $\{a, b, c\}^{\pm}$.

If a finite graph Γ is not well-labelled then a process of iterative identifications of each pair $\{e_1, e_2\}$ of distinct edges with the same initial vertex and the same label to a single edge yields a well-labelled graph. Such identifications are called *foldings*, and the whole process is known as the process of *Stallings' foldings* [3, 19, 29, 30].

Thus the graph Γ_2 on Figure 1 is obtained from the graph Γ_1 by folding the edges e_1 and e_2 to a single edge labelled by a.

Notice that the graph Γ_3 is obtained from the graph Γ_2 by removing the edge labelled by a whose initial vertex has degree 1. Such an edge is called a *hair*, and the above procedure is used to be called *"cutting hairs"*.

The label of a path $p = e_1 e_2 \cdots e_n$ in Γ, where $e_i \in E(\Gamma)$, is the word

$$lab(p) \equiv lab(e_1) \cdots lab(e_n) \in (X^{\pm})^*.$$

Notice that the label of the empty path is the empty word. As usual, we identify the word $lab(p)$ with the corresponding element in $G = gp\langle X|R\rangle$. We say that p is a *normal path* (or p is a path in *normal form*) if $lab(p)$ is a normal word.

If Γ is a well-labelled graph then a path p in Γ is freely reduced if and only if $lab(p)$ is a freely reduced word. Otherwise p can be converted into a freely reduced path p' by iterative removals of the subpaths $e\overline{e}$ (*backtrackings*) ([29, 19]). Thus

$$\iota(p') = \iota(p), \;\; \tau(p') = \tau(p) \;\; \text{and} \;\; lab(p) =_{FG(X)} lab(p'),$$

where $\boxed{FG(X)}$ is a free group with a free basis X. We say that p' is obtained from p by *free reductions*.

If $v_1, v_2 \in V(\Gamma)$ and p is a path in Γ such that

$$\iota(p) = v_1, \;\; \tau(p) = v_2 \text{ and } lab(p) \equiv u,$$

then, following the automata theoretic notation, we simply write $\boxed{v_1 \cdot u = v_2}$ to summarize this situation, and say that the word u is *readable* at v_1 in Γ.

A pair $\boxed{(\Gamma, v_0)}$ consisting of the graph Γ and the *basepoint* v_0 (a distinguished vertex of the graph Γ) is called a *pointed graph*.

Following the notation of Gitik ([11]) we denote the set of all closed paths in Γ starting at v_0 by $\boxed{Loop(\Gamma, v_0)}$, and the image of $lab(Loop(\Gamma, v_0))$ in $G = gp\langle X|R\rangle$ by $\boxed{Lab(\Gamma, v_0)}$. More precisely,

$$Loop(\Gamma, v_0) = \{p \mid p \text{ is a path in } \Gamma \text{ with } \iota(p) = \tau(p) = v_0\},$$

$$Lab(\Gamma, v_0) = \{g \in G \mid \exists p \in Loop(\Gamma, v_0) \ : \ lab(p) =_G g\}.$$

It is easy to see that $Lab(\Gamma, v_0)$ is a subgroup of G ([11]). Moreover, $Lab(\Gamma, v) = gLab(\Gamma, u)g^{-1}$, where $g =_G lab(p)$, and p is a path in Γ from v to u ([19]). If $V(\Gamma) = \{v_0\}$ and $E(\Gamma) = \emptyset$ then we assume that $H = \{1\}$.

We say that $H = Lab(\Gamma, v_0)$ is *the subgroup of G determined by the graph* (Γ, v_0). Thus any pointed graph labelled by $X^{\pm}$, where X is a generating set of a group G, determines a subgroup of G. This argues the use of the name *subgroup graphs* for such graphs.

Morphisms of Labelled Graphs. Let Γ and Δ be graphs labelled with $X^{\pm}$. The map $\pi : \Gamma \to \Delta$ is called a *morphism of labelled graphs*, if π takes vertices to vertices, edges to edges, preserves labels of direct edges and has the property that

$$\iota(\pi(e)) = \pi(\iota(e)) \text{ and } \tau(\pi(e)) = \pi(\tau(e)), \ \forall e \in E(\Gamma).$$

An injective morphism of labelled graphs is called an *embedding*. If π is an embedding then we say that the graph Γ *embeds* in the graph Δ.

A *morphism of pointed labelled graphs* $\pi : (\Gamma_1, v_1) \to (\Gamma_2, v_2)$ is a morphism of underlying labelled graphs $\pi : \Gamma_1 \to \Gamma_2$ which preserves the basepoint $\pi(v_1) = v_2$. If Γ_2 is well-labelled then there exists at most one such morphism ([19]).

Remark 3.1 ([19])**.** If two pointed well-labelled (with $X^{\pm}$) graphs (Γ_1, v_1) and (Γ_2, v_2) are isomorphic, then there exists a unique isomorphism $\pi : (\Gamma_1, v_1) \to (\Gamma_2, v_2)$. Therefore (Γ_1, v_1) and (Γ_2, v_2) can be identified via π. In this case we sometimes write $(\Gamma_1, v_1) = (\Gamma_2, v_2)$. ◇

The notation $\Gamma_1 = \Gamma_2$ means that there exists an isomorphism between these two graphs. More precisely, one can find $v_i \in V(\Gamma_i)$ $(i \in \{1, 2\})$ such that $(\Gamma_1, v_1) = (\Gamma_2, v_2)$ in the sense of Remark 3.1.

Lemma 3.2 ([19])**.** *Let (Γ_1, v_1) and (Γ_2, v_2) be pointed graphs well-labelled with $X^{\pm}$ such that $degree(v) > 1$* [2] *, for all $v \in V(\Gamma_i) \setminus \{v_i\}$ $(i \leq \{1, 2\})$.*

Then $Lab(\Gamma_1, v_1) \leq Lab(\Gamma_2, v_2)$ if and only if there exists a unique morphism $\pi : (\Gamma_1, v_1) \to (\Gamma_2, v_2)$. ◇

---

[2] Recall $degree(v) = |\{e \in E(\Gamma_i) \mid \iota(e) = v \text{ or } \tau(e) = v\}|$.

4. Subgroup Graphs

The current section is devoted to the discussion on subgroup graphs constructed by the generalized Stallings' folding algorithm. The main results of [32] concerning these graphs (more precisely, Theorem 7.1, Lemma 8.6, Lemma 8.7, Theorem 8.9 and Corollary 8.11 in [32]), which are essential for the present paper, are summarized in Theorem 4.1 below. All the missing notations are explained along the rest of the present section.

Theorem 4.1. *Let* $H = \langle h_1, \cdots, h_k \rangle$ *be a finitely generated subgroup of an amalgam of finite groups* $G = G_1 *_A G_2$.

Then there is an algorithm (the generalized Stallings' folding algorithm) which constructs a finite labelled graph $(\Gamma(H), v_0)$ *with the following properties:*

1. $Lab(\Gamma(H), v_0) = H$.
2. *Up to isomorphism,* $(\Gamma(H), v_0)$ *is a unique reduced precover of* G *determining* H.
3. $(\Gamma(H), v_0)$ *is the normal core of* $(Cayley(G, H), H \cdot 1)$.
4. *A normal word* $g \in G$ *is in* H *if and only if it labels a closed path in* $\Gamma(H)$ *starting at* v_0*, that is* $v_0 \cdot g = v_0$.
5. *Let* m *be the sum of the lengths of words* $h_1, \dots h_n$. *Then the algorithm computes* $(\Gamma(H), v_0)$ *in time* $O(m^2)$. *Moreover,* $|V(\Gamma(H))|$ *and* $|E(\Gamma(H))|$ *are proportional to* m.

Corollary 4.2. *Theorem 4.1 (4) provides a solution of the membership problem for finitely generated subgroups of amalgams of finite groups.*

Throughout the present paper the notation $\boxed{(\Gamma(H), v_0)}$ is always used for the finite labelled graph constructed by the generalized Stallings' folding algorithm for a finitely generated subgroup H of an amalgam of finite groups $G = G_1 *_A G_2$.

Definition of Precovers: The notion of *precovers* was defined by Gitik in [11] for subgroup graphs of amalgams. Below we present its definition and list some basic properties. In doing so, we rely on the notation and results obtained in [11]. The discussion of precovers which are *reduced* come later in Section 5.

Let Γ be a graph labelled with $X^{\pm}$, where $X = X_1 \cup X_2$ is the generating set of $G = G_1 *_A G_2$ given by (1.a)-(1.c). We view Γ as a two colored graph: one color for each one of the generating sets X_1 and X_2 of the factors G_1 and G_2, respectively.

The vertex $v \in V(\Gamma)$ is called X_i*-monochromatic* if all the edges of Γ incident with v are labelled with $X_i^{\pm}$, for some $i \in \{1, 2\}$. We denote the set of X_i-monochromatic vertices of Γ by $VM_i(\Gamma)$ and put $VM(\Gamma) = VM_1(\Gamma) \cup VM_2(\Gamma)$.

We say that a vertex $v \in V(\Gamma)$ is *bichromatic* if there exist edges e_1 and e_2 in Γ with

$$\iota(e_1) = \iota(e_2) = v \text{ and } lab(e_i) \in X_i^{\pm}, \ i \in \{1, 2\}.$$

The set of bichromatic vertices of Γ is denoted by $VB(\Gamma)$.

A subgraph of Γ is called *monochromatic* if it is labelled only with $X_1^{\pm}$ or only with $X_2^{\pm}$. An *X_i-monochromatic component* of Γ ($i \in \{1, 2\}$) is a maximal connected subgraph of Γ labelled with $X_i^{\pm}$, which contains at least one edge. Thus monochromatic components of Γ are graphs determining subgroups of the factors, G_1 or G_2.

We say that a graph Γ is *G-based* if any path $p \subseteq \Gamma$ with $lab(p) =_G 1$ is closed. Thus if Γ is G-based then, obviously, it is well-labelled with $X^{\pm}$.

Definition 4.3 (Definition of Precover)**.** *A G-based graph Γ is a* precover *of G if each X_i-monochromatic component of Γ is a* cover *of G_i ($i \in \{1, 2\}$).*

Following the terminology of Gitik ([11]), we use the term *"covers of G"* for *relative (coset) Cayley graphs* of G and denote by $\boxed{Cayley(G, S)}$ the coset Cayley graph of G relative to the subgroup S of G.[3] If $S = \{1\}$, then $Cayley(G, S)$ is the *Cayley graph* of G and the notation $\boxed{Cayley(G)}$ is used.

Note that the use of the term "covers" is adjusted by the well known fact that a geometric realization of a coset Cayley graph of G relative to some $S \leq G$ is a 1-skeleton of a topological cover corresponding to S of the standard 2-complex representing the group G (see [47], pp.162-163).

Convention 4.4. *By the above definition, a precover doesn't have to be a connected graph. However along this paper we restrict our attention only to connected precovers. Thus any time this term is used, we always mean that the corresponding graph is connected unless it is stated otherwise.*

We follow the convention that a graph Γ with $V(\Gamma) = \{v\}$ and $E(\Gamma) = \emptyset$ determining the trivial subgroup (that is $Lab(\Gamma, v) = \{1\}$) is a (an empty) precover of G. ◇

Example 4.5. Let $G = gp\langle x, y | x^4, y^6, x^2 = y^3 \rangle = \mathbb{Z}_4 *_{\mathbb{Z}_2} \mathbb{Z}_6$.

Recall that G is isomorphic to $SL(2, \mathbb{Z})$ under the homomorphism

$$x \mapsto \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \ y \mapsto \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}.$$

---

[3] Whenever the notation $Cayley(G, S)$ is used, it always means that S is a subgroup of the group G and the presentation of G is fixed and clear from the context.

The graphs Γ_1 and Γ_3 on Figure 2 are examples of precovers of G with one monochromatic component and two monochromatic components, respectively.

Though the $\{x\}$-monochromatic component of the graph Γ_2 is a cover of $\mathbb{Z}_4$ and the $\{y\}$-monochromatic component is a cover of $\mathbb{Z}_6$, Γ_2 is not a precover of G, because it is not a G-based graph. Indeed, $v \cdot (x^2y^{-3}) = u$, while $x^2y^{-3} =_G 1$.

The graph Γ_4 is not a precover of G because its $\{x\}$-monochromatic components are not covers of $\mathbb{Z}_4$. ◇



FIGURE 2.

A graph Γ is *x-saturated* at $v \in V(\Gamma)$, if there exists $e \in E(\Gamma)$ with $\iota(e) = v$ and $lab(e) = x$ $(x \in X)$. Γ is *$X^{\pm}$-saturated* if it is x-saturated for each $x \in X^{\pm}$ at each $v \in V(\Gamma)$.

Lemma 4.6 (Lemma 1.5 in [11]). *Let $G = gp\langle X|R\rangle$ be a group and let (Γ, v_0) be a graph well-labelled with $X^{\pm}$. Denote $Lab(\Gamma, v_0) = S$. Then*

- *Γ is G-based if and only if it can be embedded in $(Cayley(G,S), S\cdot 1)$,*
- *Γ is G-based and $X^{\pm}$-saturated if and only if it is isomorphic to $(Cayley(G,S), S\cdot\ 1)$.* [4]

Corollary 4.7. *If Γ is a precover of G with $Lab(\Gamma, v_0) = H \leq G$ then Γ is a subgraph of $Cayley(G,H)$.*

Thus a precover of G can be viewed as a part of the corresponding cover of G, which explains the use of the term "precovers".

Remark 4.8 ([32]). Let $\phi : \Gamma \to \Delta$ be a morphism of labelled graphs. If Γ is a precover of G, then $\phi(\Gamma)$ is a precover of G as well. ◇

---

[4] We write $S\cdot 1$ instead of the usual $S1 = S$ to distinguish this vertex of $Cayley(G,S)$ as the basepoint of the graph.

Precovers are Compatible: A graph Γ is called *compatible at a bichromatic vertex* v if for any monochromatic path p in Γ such that $\iota(p) = v$ and $lab(p) \in A$ there exists a monochromatic path t of a different color in Γ such that $\iota(t) = v$, $\tau(t) = \tau(p)$ and $lab(t) =_G lab(p)$. We say that Γ is *compatible* if it is compatible at all bichromatic vertices.

Example 4.9. The graphs Γ_1 and Γ_3 on Figure 2 are compatible. The graph Γ_2 does not possess this property because $w \cdot x^2 = v$, while $w \cdot y^3 = u$. Γ_4 is not compatible as well. ◇

Lemma 4.10 (Lemma 2.12 in [11])**.** *If Γ is a compatible graph, then for any path p in Γ there exists a path t in normal form such that $\iota(t) = \iota(p)$, $\tau(t) = \tau(p)$* and *$lab(t) =_G lab(p)$.*

Remark 4.11 (Remark 2.11 in [11])**.** Precovers are compatible. ◇

The following can be taken as another definition of precovers.

Lemma 4.12 (Corollary2.13 in [11])**.** *Let Γ be a compatible graph. If all X_i-components of Γ are G_i-based, $i \in \{1, 2\}$, then Γ is G-based. In particular, if each X_i-component of Γ is a cover of G_i, $i \in \{1, 2\}$, and Γ is compatible, then Γ is a precover of G.*

Normal Core and Canonicity:

Definition 4.13. *A vertex of $Cayley(G, H)$ is called essential if there exists a normal path closed at $H \cdot 1$ that goes through it.*

The normal core $(\Delta, H{\cdot}1)$ of $Cayley(G, H)$ is the restriction of $Cayley(G, H)$ to the set of all essential vertices.

Remark 4.14. Note that the normal core $(\Delta, H \cdot 1)$ can be viewed as the union of all normal paths closed at $H \cdot 1$ in $(Cayley(G, H), H \cdot 1)$. Thus $(\Delta, H \cdot 1)$ is a connected graph with basepoint $H \cdot 1$.

Moreover, $V(\Delta) = \{H \cdot 1\}$ and $E(\Delta) = \emptyset$ if and only if H is the trivial subgroup. Indeed, H is not trivial iff there exists $1 \neq g \in H$ in normal form iff g labels a normal path in $Cayley(G, H)$ closed at $H \cdot 1$, iff $E(\Delta) \neq \emptyset$.

◇

Therefore the normal core of $Cayley(G, H)$ depends on H itself and not on the set of subgroup generators, which, by Theorem 4.1 (3), implies the canonicity of the construction of $(\Gamma(H), v_0)$ by the generalized Stallings' folding algorithm. This provides a solution of the Membership Problem for finitely generated subgroups of amalgams of finite groups given by Theorem 4.1 (4).

Complexity Issues: As were noted in [32], the complexity of the generalized Stallings' algorithm is quadratic in the size of the input, when we assume that all the information concerning the finite groups G_1, G_2, A and the amalgam $G = G_1 *_A G_2$ given via $(1.a)$, $(1.b)$ and $(1.c)$ (see Section 3) is not a part of the input. We also assume that the Cayley graphs and all the relative Cayley graphs of the free factors are given for "free" as well.

Otherwise, if the group presentations of the free factors G_1 and G_2, as well as the monomorphisms between the amalgamated subgroup A and the free factors are a part of the input (the *uniform version* of the algorithm) then we have to build the groups G_1 and G_2, that is to construct their Cayley graphs and relative Cayley graphs.

Since we assume that the groups G_1 and G_2 are finite, the Todd-Coxeter algorithm and the Knuth Bendix algorithm are suitable [27, 45, 47] for these purposes. Then the complexity of the construction depends on the group presentation of G_1 and G_2 we have: it could be even exponential in the size of the presentation [7]. Therefore the generalized Stallings algorithm, presented in [32], with these additional constructions could take time exponential in the size of the input.

Thus each uniform algorithmic problem for H whose solution involves the construction of the subgroup graph $\Gamma(H)$ may have an exponential complexity in the size of the input.

The primary goal of the complexity analysis introduced along the current paper is to estimate our graph theoretical methods. To this end, we assume that all the algorithms along the present paper have the following "given data".

GIVEN: : Finite groups G_1, G_2, A and the amalgam $G = G_1 *_A G_2$ given via $(1.a)$, $(1.b)$ and $(1.c)$.
We assume that the Cayley graphs and all the relative Cayley graphs of the free factors are given.

5. The Conjugacy Problem

The *conjugacy problem* for subgroups of a group G asks to answer whether or not given subgroups of G are conjugate. Below we solve this problem for finitely generated subgroups of amalgams of finite groups, using subgroup graphs constructed by the generalized Stallings' algorithm.

Our results extend the analogous ones obtained for finitely generated subgroups of free groups by Kapovich and Myasnikov in [19]. We start by discussing of this analogy. Throughout the present section we assume that $G = G_1 *_A G_2$ is an amalgam of finite groups.

The solution of the *conjugacy problem* for finitely generated subgroups of free groups, presented in [19], involve a construction of a special graph $Type$ which is a *core graph* with respect to each of its vertices. Thus it posses the property that $H, K \leq_{f.g.} FG(X)$ conjugate if and only if $Type(\Gamma_H) = Type(\Gamma_K)$.

The extended definition of $Type$ in the case of amalgams of finite groups as well as a discussion of its properties are introduced in Section 5.1. Theorem 5.16 gives a connection between $Types$ of conjugate subgroups, which provides a solution of the conjugacy problem for subgroups in amalgams of finite groups. The algorithm is presented along with the proof of Corollary 5.17. The complexity analysis shows that this algorithm is quadratic in the size of the input.

In [46] Stallings defined a *core-graph* to be a connected graph which has at least one edge, and each of whose edges belongs to at least one cyclically reduced circuit. He noted that every connected graph with a non-trivial fundamental group contains a *core* where the fundamental group is concentrated, and the original graph consists of this core with various trees *hangings* on. Thus given a connected graph Γ which has at least one edge, one can obtain its core by the process of "*shaving off trees*".

In [19] the Stallings' notion of a *core-graph* were split into two aspects: a core with respect to some vertex (*the basepoint*) and a core with respect to any of its vertices. The first notion corresponds to the subgroup graph (Γ_S, v_0) of $S \leq_{f.g.} F(X)$ constructed by Stallings' algorithm [46], while the second one defines $Type(\Gamma_S)$. Thus Γ_S can be obtained from $Cayley(FG(X), S)$ by a "partial shaving procedure", which preserves the basepoint $S \cdot 1$. The "full shaving procedure" yields $Type(\Gamma_S)$. Moreover, $Type(\Gamma_S)$ can be obtained from the subgroup Γ_S by the iterative erasure of the unique sequence of *spurs* (*spur* is an edge one of whose endpoints has degree 1) starting from the basepoint v_0 of Γ_S.

An analog of (Γ_S, v_0) in amalgams of finite groups is the subgroup graph $(\Gamma(H), v_0)$ constructed by the generalized Stallings' algorithm, where $H \leq_{f.g.} G_1 *_A G_2$. By Theorem 4.1 (3), $(\Gamma(H), v_0)$ is the *normal core* of $(Cayley(G, H), H \cdot 1)$, that is the union of all normal paths in $(Cayley(G, H), H \cdot 1)$ closed at $H \cdot 1$. That is, it is a sort of a core graph with respect to the basepoint $H \cdot 1$.

An analog of a *spur* in subgroup graphs of finitely generated subgroups of amalgams of finite groups is a *redundant component.* The notion of *redundant component* were defined in [32]. However in the present context its more convenient to use the name *redundant component w.r.t.* the basepoint v_0 for that notion defined in [32], and to keep the name *redundant component* for the following.

Definition 5.1. *Let* Γ *be a precover of* G. *Let* C *be a* X_i*-monochromatic component of* Γ $(i \in \{1,2\})$. C *is* *redundant* *if one of the following holds.*

1. C *is the unique monochromatic component of* Γ *(that is* $\Gamma = C$*) and* $Lab(C,v) = \{1\}$ *(equivalently, by Lemma 4.6,* C *is isomorphic to* $Cayley(G_i)$*), where* $v \in V(C)$.
2. Γ *has at least two distinct monochromatic components and the following holds.*

   Let $\vartheta \in VB(C)$. *Let* $K = Lab(C,\vartheta)$ *(equivalently, by Lemma 4.6,* $(C,\vartheta) = (Cayley(G_i,K), K \cdot 1)$*).*

   Then $K \leq A$ *and* $VB(C) = A(\vartheta)$. [5]

C *is* *redundant w.r.t.* *the vertex* $u \in V(\Gamma)$ *if* C *is redundant and* $u \in V(C)$ *implies* $u \in VB(C)$ *and* $K = \{1\}$.

Remark 5.2. Similarly to the removing of spurs from graphs representing finitely generated subgroups of free group, in the case of amalgams of finite groups the erasing of redundant components w.r.t. v_0 from (Γ, v_0) doesn't change the subgroup defined by this pointed graph (see Lemma 6.17 in [32]).

Namely, if Γ' is the graph obtained from Γ, by erasing of a monochromatic component which is redundant w.r.t. v_0, then $Lab(\Gamma', v_0') = Lab(\Gamma, v_0)$, where v_0' is the image of v_0 in Γ'.

◇

The following example attempts to give an intuition of what happens in the covering space corresponding to the subgroup $H \leq G$ of the standard 2-complex representing G, when we remove redundant monochromatic components from a subgraph of $Cayley(G,H)$, which is the 1-skeleton of this covering space.

Example 5.3. Let $G = gp\langle x, y | x^4, y^6, x^2 = y^3 \rangle = G_1 *_A G_2$, where $G_1 = gp\langle x | x^4 \rangle$, $G_2 = gp\langle y | y^6 \rangle$ and $A = \langle x^2 \rangle = \langle y^3 \rangle$.

Assume that all the redundant monochromatic components are isomorphic to either $Cayley(G_1)$ or $Cayley(G_2)$. Hence a removing of a redundant component from $Cayley(G,H)$ is expressed in the covering space by removing a 2-cell with the boundary path x^4 (or y^6) and two 2-cells with the boundary path x^2y^{-3}. One can imagine this process as "smashing of bubbles", see Figure 3.

However even if a redundant component is isomorphic to $Cayley(G_i, S)$, where $\{1\} \neq S \leq G_i$, $i \in \{1,2\}$, the "bubbles intuition" fails as well as in the cases when the factor groups are not cyclic. That is now the parts removed from the covering space hardly resemble bubbles, while the motivation for their removing remains the same.

---

[5] Recall that $A(\vartheta) = \{\vartheta \cdot a \mid a \in A\}$ is the *A-orbit* of ϑ in $V(C)$ by the right action of A on $V(C)$. Since $A_\vartheta = K$, the condition $VB(C) = A(\vartheta)$ can be replaced by its computational analogue $|VB(C)| = [A : K]$.

Here the common intuition with Stallings' construction: we "smash bubbles" instead of "shaving off trees", which can be thought of as an iterative erasure of spurs. ◇



FIGURE 3. A bubble.

The subgroup graph $(\Gamma(H), v_0)$ is a unique finite *reduced precover* of G, by Theorem 4.1 (2). Now we are ready to recall the precise definition of this term.

Definition 5.4 (Definition 6.18 in [32]). *A precover* (Γ, v_0) *of* G *is* *reduced* *if the following holds.*

(i) (Γ, v_0) *has no redundant components w.r.t.* v_0*.*
(ii) $Lab(C_0, v_0) \cap A \neq \{1\}$ *implies* $v_0 \in VB(\Gamma)$*, where* C_0 *is a monochromatic component of* Γ *such that* $v_0 \in V(C_0)$*.*

Roughly speaking, the reduced precover $(\Gamma(H), v_0)$ can be obtained from $(Cayley(G, H), H \cdot 1)$ by removing of all redundant components w.r.t. the basepoint $H \cdot 1$. Intuitively, in analogy with [19], the graph obtained from $(Cayley(G, H), H \cdot 1)$ by erasing of all redundant components is $Type(\Gamma(H))$. Moreover, $Type(\Gamma(H))$ can be obtained from the graph $\Gamma(H)$ by the iterative erasure of the unique sequence of redundant components starting from C_0 such that $v_0 \in V(C_0)$. Some special cases occur when H is a subgroup of a factor, G_1 or G_2, of G.

5.1. **Type.**

Consider $(\Gamma(H), v_0)$, where H is a finitely generated subgroup of an amalgam of finite groups $G = G_1 *_A G_2$.

As is mentioned in the introductory part, a definition of $Type(\Gamma(H))$ largely relies on the definition of $\Gamma(H)$. To this end we start by presenting some properties of reduced precovers based on the results obtained in [32].

Lemma 5.5 (Lemma 6.21 in [32])**.** *Let* (Γ, v) *be a precover of* G *with no redundant components w.r.t.* v*. Let* $H = Lab(\Gamma, v)$*.*

If (Γ, v) *is not a reduced precover of* G*, then* $Lab(C, v) \cap A = S \neq \{1\}$ *and* $v \in VM_i(\Gamma)$*, where* C *is a* X_i*-monochromatic component of* Γ *such that* $v \in V(C)$ *(*$i \in \{1, 2\}$*).*

Moreover, $(\Gamma(H), v_0) = (\Gamma *_{\{v \cdot a = Sa \mid a \in A\}} Cayley(G_j, S), \vartheta)$*,where* $1 \leq i \neq j \leq 2$ *and* ϑ *is the image of* v *(equivalently, of* $S \cdot 1$*) in the amalgam graph.*

Corollary 5.6. *Let* Γ *be a precover of* G*. Let* C *be a* X_i*-monochromatic component of* Γ *and let* $v \in VM_i(C)$ *(*$i \in \{1, 2\}$*).*

Then the graph $\Delta = \Gamma *_{\{v \cdot a = Sa \mid a \in A\}} Cayley(G_j, S)$*, where* $S = Lab(C, v) \cap A$*, satisfies*

- Γ *and* $Cayley(G_j, S)$ *embeds in* Δ *(*$1 \leq i \neq j \leq 2$*),*
- $Lab(\Delta, \vartheta) = Lab(\Gamma, v)$*, where* ϑ *is the image of* v *in* Δ*.*

Lemma 5.7. *Each of the following holds.*

(i) $\underline{H = \{1\}}$ *if and only if* $V(\Gamma(H)) = \{v_0\}$*,* $E(\Gamma(H)) = \emptyset$*.*

(ii) $\underline{H \leq G_i \text{ and } H \cap A = \{1\}}$ *if and only if* $\Gamma(H)$ *consists of a unique* X_i*-monochromatic component:* $(\Gamma(H), v_0) = (Cayley(G_i, H), H \cdot 1)$ *(*$i \in \{1, 2\}$*).*

(iii) $\underline{H \leq A}$ *if and only if* $(\Gamma(H), v_0) = (\Delta, \vartheta)$*, where* $\Delta = Cayley(G_1, H) *_{\{Ha \mid a \in A\}} Cayley(G_2, H)$ *and* ϑ *is the image of* $H \cdot 1$ *in* Δ*.*

(iv) *If* $\underline{H \not\leq G_i \text{ for all } i \in \{1, 2\}}$ *then* $C_0 \subseteq \Gamma(H)$ *is a redundant component if and only if* $v_0 \in V(C_0)$ *and there exists* $u_0 \in VB(C_0)$ *such that* $VB(C_0) = A(u_0)$ *and* $Lab(C_0, u_0) \leq A$*. Moreover,* C_0 *is a unique redundant component of* $\Gamma(H)$*.*

Proof. By Theorem 4.1 (2), the reduced precover $(\Gamma(H), v_0)$ is unique up to isomorphism. By the Definition 5.4, a graph (Δ, u) such that $V(\Delta) = \{u\}$ and $E(\Delta) = \emptyset$ is a reduced precover of G, which satisfies $Lab(\Delta, u) = \{1\}$. Therefore $(\Delta, u) = (\Gamma(H), v_0)$. This gives the "if" direction of (i). Similar arguments prove the "if" direction of (ii) and (iii).

The converse of (i) is trivial. The opposite direction of (ii) is true, because, by Lemma 4.6, $(\Gamma, v) = (Cayley(G_i, H), H \cdot 1)$ ($i \in \{1, 2\}$) implies $Lab(\Gamma, v) = H$. Moreover, by Definition 5.4, (Γ, v) is a reduced precover of G with $Lab(\Gamma, v) = H$ whenever $H \leq G_i$ such that $H \cap A = \{1\}$.

To prove the converse of (iii), let $\Gamma = C_1 *_{\{v_1 \cdot a = v_2 \cdot a \mid a \in A\}} C_2$, where $(C_i, v_i) = (Cayley(G_i, H), H \cdot 1)$ and $H \leq A$ ($i \in \{1, 2\}$). Let v be the image of v_i in Γ. Hence $Lab(C_i, v_i) \leq Lab(\Gamma, v)$. Now we need the following result from [32].

Claim 1. *Let* (Γ, v) *be a precover of* G. *Let* C *be a* X_i*-monochromatic component of* Γ. *Then the followings are equivalent.*

- $u_1 \cdot a = u_2$ *implies* $a \in A$, *for all* $u_1, u_2 \in VB(C)$.
- $VB(C) = A(\vartheta)$ *and* $Lab(C, \vartheta) \leq A$, *for all* $\vartheta \in VB(C)$.

Thus $u_1 \cdot a = u_2$ implies $a \in A$, for all $u_1, u_2 \in VB(C_1) = VB(C_2)$. Therefore no normal words of syllable length greater than 1 label normal paths in Γ closed at v. Hence if $g \in Lab(\Gamma, v)$ and p is a normal path in Γ closed at v such that $lab(p) \equiv g$ then either $p \subseteq C_1$ or $p \subseteq C_2$. Thus $g \equiv lab(p) \in H$. Therefore $Lab(\Gamma, v) = H \leq A$. By Theorem 4.1 (2), $(\Gamma, v) = (\Gamma(H), v_0)$.

The statement of (iv) is an immediate consequence of Definition 5.4 and Definition 5.1.

To prove the uniqueness of C_0 assume that there exists another redundant component D in $\Gamma(H)$ such that $v_0 \in V(D)$ and there exists $u \in VB(D)$ such that $VB(D) = A(u)$ and $Lab(D, u) \leq A$. Thus, without loss of generality, one can assume that C_0 is a X_1-monochromatic component and D is a X_2-monochromatic component. Hence $v_0 \in VB(C_0) \cap VB(D)$. Therefore $A(u_0) = A(v_0) = A(u)$. Hence $VB(C_0) = VB(D)$. Since the graph $\Gamma(H)$ is well-labelled, this implies that C_0 and D are the only monochromatic components of $\Gamma(H)$.

Moreover, since $v_0 \in A(u_0)$, there is $a \in A$ such that $v_0 = u_0 \cdot a$. Hence $Lab(C_0, v_0) = aLab(C_0, u_0)a^{-1} \leq A$. Similarly, $Lab(D, v_0) \leq A$. Thus $Lab(C_0, v_0) = A_{v_0} = Lab(D, v_0)$. [6] Therefore, by (iii), $H = A_{v_0} \leq A$, which contradicts the assumption of (iv).

◇

Remark 5.8. In (iii), the graphs $Cayley(G_1, H)$ and $Cayley(G_2, H)$ embeds in Δ, by Corollary 5.6, ◇

Corollary 5.9 (The Triviality Problem)**.** *Let* $h_1, \cdots, h_n \in G$.

Then there is an algorithm which decides whether or not the subgroup $H = \langle h_1, \cdots, h_n \rangle$ *is trivial.*

Proof. We first construct the pointed graph $(\Gamma(H), v_0)$, using the generalized Stallings' folding algorithm.

By Theorem 4.1 (2), $(\Gamma(H), v_0)$ is a reduced precover of G. Therefore, by Lemma 5.7(i), $H = Lab(\Gamma(H), v_0) = \{1\}$ if and only if $V(\Gamma(H)) = v_0$ and $E(\Gamma(H)) = \emptyset$. ◇

Remark 5.10 (Complexity)**.** To detect the triviality of a subgroup H given by a set of generators it takes the same time as to construct the subgroup graph $\Gamma(H)$. By Theorem 4.1 (5), it is $O(m^2)$, where m is the sum of the lengths of words $h_1, \dots h_n$. ◇

---

[6] $A_{v_0} = Lab(C_0, v_0) \cap A$ is the A-*stabilizer* of v_0 by the right action of A on $V(C_0)$.

Lemma 5.11. *Let H be a finitely generated subgroup of G such that $H \nleq G_i$ ($i \in \{1,2\}$).*

If $\Gamma(H)$ has a redundant component, then there exists a unique sequence of alternating monochromatic components $C_0, \cdots C_{m-1}$ of $\Gamma(H)$ such that the graph Γ_m, obtained from $\Gamma(H)$ by the iterative erasure of the above sequence, has no redundant components.

Proof. By Lemma 5.7 (iv), $\Gamma(H)$ has the unique redundant component C_0 which satisfies $v_0 \in V(C_0)$ and there exists $u_0 \in VB(C_0)$ such that $VB(C_0) = A(u_0)$ and $Lab(C_0, u_0) \leq A$.

Let Γ_1 be the graph obtained from $\Gamma(H)$ by removing of the component C_0. That is

$$VM_i(\Gamma_1) = VM_i(\Gamma(H)) \setminus VM_i(C_0), \quad VM_j(\Gamma_1) = VM_j(\Gamma(H)),$$

$$VB(\Gamma_1) = VB(\Gamma(H)) \setminus VB(C_0) \text{ and } E(\Gamma_1) = E(\Gamma(H)) \setminus E(C_0).$$

The resulting graph Γ_1 is, obviously, a finite precover of G. If Γ_1 has no redundant components then $m = 1$.

Otherwise there exists a unique X_j-monochromatic component of Γ_1 ($1 \leq i \neq j \leq 2$) which is redundant. Indeed, $\Gamma(H)$ has a unique X_j-monochromatic component C_1 such that $C_0 \cap C_1 = VB(C_0)$ ($1 \leq i \neq j \leq 2$). By abuse of notation, we identify the component C_1 of $\Gamma(H)$ with its image in Γ_1. Thus

$$VB_{\Gamma_1}(C_1) = VB_{\Gamma(H)}(C_1) \setminus VB(C_0).$$

Therefore, C_1 is a X_j-monochromatic redundant component of Γ_1 if and only if there exists a vertex $u_1 \in VB_{\Gamma(H)}(C_1) \setminus VB(C_0)$ such that $Lab(C_1, u_1) \leq A$ and $VB_{\Gamma(H)}(C_1) = A(u_1) \cup VB(C_0) = A(u_1) \cup A(u_0)$.

Since the graph $\Gamma(H)$ is finite, continuing in such manner one can find the unique sequence

$$C_0, C_1, \dots, C_{m-1} \tag{$*$}$$

of X_{j_i}-monochromatic components of $\Gamma(H)$ (see Figure 4) such that the following holds.

1. $1 \leq j_i \neq j_{i+1} \leq 2$ for all $0 \leq i \leq m-1$.
2. $v_0 \in V(C_0)$ and there exists $u_0 \in VB(C_0)$ such that $Lab(C_0, u_0) \leq A$ and $VB(C_0) = A(u_0)$.
3. For all $1 \leq i \leq m-1$, there exists $u_i \in VB(C_i) \setminus VB(C_{i-1})$ such that $Lab(C_i, u_i) \leq A$ and $VB(C_i) = A(u_{i-1}) \cup A(u_i)$.
4. The graph Γ_m, obtained from $\Gamma(H)$ by the iterative removal of sequence $(*)$, has no redundant components.

◇

Following the notation of Lemma 5.11 we define.

Definition 5.12 (Definition of Type)**.** *Let H be a finitely generated subgroup of $G = G_1 *_A G_2$.*

If $H \leq G_i$ or $\Gamma(H)$ has no redundant components then $Type(\Gamma(H)) = \Gamma(H)$. Otherwise $Type(\Gamma(H)) = \Gamma_m$.

Lemma 5.13 (Properties of $Type(\Gamma(H))$)**.**

(i) *$Type(\Gamma(H))$ is a finite nonempty precover of G.*

Let $v \in V(Type(\Gamma(H)))$. Let $K = Lab(Type(\Gamma(H)), v)$.

(ii) *$K \neq \{1\}$.*

(iii) *$Lab(\Gamma(H), v) = K$.*

(iv) *If $H, K \not\leq A$ then $Type(\Gamma(K)) = Type(\Gamma(H))$.*
If $H \leq A$ then $Type(\Gamma(H)) = C_1 *_{\{v_1 \cdot a = v_2 \cdot a \mid a \in A\}} C_2$ and

$$Type(\Gamma(K)) = \begin{cases} C_l, & K \not\leq A; \\ C_l *_{\{v \cdot a = K \cdot a \mid a \in A\}} Cayley(G_j, K), & K \leq A, \end{cases}$$



FIGURE 4. Let $H \leq_{f.g} G_1 *_A G_2 \simeq Z_4 *_{Z_2} Z_6$, where $G_1 = gp\langle x | x^4 \rangle$, $G_2 = gp\langle y | y^6 \rangle$ and $A = \langle x^2 \rangle = \langle y^3 \rangle$.
Thus C_0, C_1, C_2 is the unique sequence of alternating monochromatic components in the graph $\Gamma(H)$ such that Γ_3 has no redundant components.
In $\Gamma(H)$: $(C_0, u_0) = Cayley(G_1)$ and $VB(C_0) = A(u_0)$; $(C_1, u_1) = Cayley(G_2)$ and $VB(C_1) = A(u_0) \cup A(u_1)$; $(C_2, u_2) = Cayley(G_1)$ and $VB(C_2) = A(u_1) \cup A(u_2)$; $(C_3, u_3) = Cayley(G_2)$, but $VB(C_3) = A(u_2) \cup A(u_3) \cup A(u_4)$. Thus $Type(\Gamma(H)) = \Gamma_3$.

where $(C_i, v_i) = (Cayley(G_i, H), H \cdot 1)$, *for all* $i \in \{1, 2\}$, $Lab(C_l, v) = K$ *and* $1 \leq l \neq j \leq 2$.

Proof. If $Type(\Gamma(H)) = \Gamma(H)$ then the statement of (i)-(iii) is trivial. Therefore, without loss of generality, we can assume that $Type(\Gamma(H)) = \Gamma_m$, where Γ_m is obtained from $\Gamma(H)$ by the iterative removal of the unique sequence $(*)$ of alternating monochromatic components

$$C_0, C_1, \ldots, C_{m-1}.$$

By the construction, $Type(\Gamma(H)) = \Gamma_m$ is a finite precover of G. Assume that Γ_m consists of a unique monochromatic component C_m, that is $\Gamma_m = C_m$, then C_m is not redundant. Indeed, $|VB_{\Gamma_m}(C_m)| = 0$ (see Figure 5), hence $VB_{\Gamma(H)}(C_m) = A(u_{m-1})$. Since $v_0 \notin V(C_m)$ and $(\Gamma(H), v_0)$ is a reduced precover of G, that is has no redundant components w.r.t. v_0, this is possible if and only if $\{1\} \neq Lab(C_m, u_{m-1}) \nleq A$. This completes the proof of (i) and (ii).



FIGURE 5. Example of the sequence C_0, C_1 of alternating monochromatic components in the graph $\Gamma(H)$, where $H \leq f.g. G_1 *_A G_2 \simeq S_3 *_{Z_2} S_3$, where $G_1 = gp\langle a, b \mid a^3,\ b^2,\ ab = ba^2 \rangle$, $G_2 = gp\langle x, y \mid x^3,\ y^2,\ xy = yx^2 \rangle$, and $A = \langle ab \rangle = \langle yx \rangle$. $\underline{\text{In } \Gamma(H)}$: $(C_0, u_0) = Cayley(G_1)$ and $VB(C_0) = A(u_0)$; $VB(C_1) = VB(C_0)$, $Lab(C_1, v) = \langle x \rangle \nleq A$. Thus $Type(\Gamma(H)) = \Gamma_2 = C_1$.

Since each monochromatic component C_i $(1 \leq i \leq m-1)$ is redundant in Γ_i (which is the graph obtained from $\Gamma(H)$ by the iterative removal of $C_0, \cdots, C_{i-1}$) w.r.t. some $v \in V(\Gamma_m) \subseteq V(\Gamma(H))$, we conclude, by Remark 5.2, that $Lab(\Gamma(H), v) = Lab(\Gamma_m, v)$. We get (iii).

To prove (iv), assume first that $H \leq A$. Therefore, by Definition 5.12 and by Lemma 5.7 (iii), $Type(\Gamma(H)) = \Gamma(H) = C_1 *_{\{v_1 \cdot a = v_2 \cdot a \mid a \in A\}} C_2$, where $(C_i, v_i) = (Cayley(G_i, H), H \cdot 1)$ $(i \in \{1, 2\})$. Without loss of generality, assume that $v_0 \neq v \in V(C_1)$. Therefore C_2 is redundant w.r.t. v. Hence, by Lemma 5.5,

$$\Gamma(K) = \begin{cases} C_1, & K \cap A = \{1\}; \\ C_1 *_{\{v \cdot a = S \cdot a \mid a \in A\}} Cayley(G_2, S), & K \cap A = S \neq \{1\}. \end{cases}$$

Therefore

$$Type(\Gamma(K)) = \begin{cases} C_1, & K \not\leq A; \\ C_1 *_{\{v \cdot a = K \cdot a \mid a \in A\}} Cayley(G_2, K), & K \leq A. \end{cases}$$

Assume now that $H, K \not\leq A$. Thus combining Definition 5.12 and Lemma 5.7, we conclude that $Type(\Gamma(H))$ has no redundant components. If $(Type(\Gamma(H)), v)$ is a finite reduced precover of G then, by Theorem 4.1 (2), $(Type(\Gamma(H)), v) = (\Gamma(K), u_0)$.

Otherwise, by Lemma 5.5,

$$(\Gamma(K), u_0) = Type(\Gamma(H)) *_{\{v \cdot a = S \cdot a \mid a \in A\}} Cayley(G_j, S),$$

where $S = Lab(C, v) \cap A \neq \{1\}$ and C is a X_i-monochromatic component of $Type(\Gamma(H))$ such that $v \in V(C)$ $(1 \leq i \neq j \leq 2)$. Since $K \not\leq A$, the component $D = Cayley(G_j, S)$ is redundant in $\Gamma(K)$. Therefore $Type(\Gamma(H)) = Type(\Gamma(K))$. This completes the proof.

◇

Example 5.14. Concerning the subgroups H_1 and H_2 from Example A.2 and their subgroup graphs $\Gamma(H_1)$ and $\Gamma(H_1)$ presented on Figures 12 and 13, we compute that $Type(\Gamma(H_1)) = \Gamma(H_1)$ and $Type(\Gamma(H_2)) = \Gamma(H_2)$. ◇

5.2. **Conjugate Subgroups.**

Lemma 5.15. *Let H and K be nontrivial subgroups of G such that $Type(\Gamma(H)) = Type(\Gamma(K))$. Then H is conjugate to K in G.*

Proof. Suppose that $Type(\Gamma(H)) = Type(\Gamma(K)) = \Gamma$. Let $v \in V(\Gamma) \subseteq V(\Gamma(H))$. Hence the subgroup $Lab(\Gamma(H), v)$ is conjugate to the subgroup $Lab(\Gamma(H), v_0)$. By Lemma 5.13 (iii), $Lab(\Gamma(H), v) = Lab(\Gamma, v)$. Therefore the subgroup $Lab(\Gamma, v)$ is conjugate to the subgroup $Lab(\Gamma(H), v_0) = H$. By symmetric arguments, the subgroup $Lab(\Gamma, v)$ is also conjugate to the subgroup K. Hence H is conjugate to K. See Figure 6. ◇

Theorem 5.16. *Let H and K be finitely generated subgroup of an amalgam of finite groups $G = G_1 *_A G_2$.*
Then H is conjugate to K in G if and only if one of the following holds

(1) *$H, K \not\leq A$ and $Type(\Gamma(K)) = Type(\Gamma(H))$.*



FIGURE 6. $K = gHg^{-1}$, where $g \equiv g_1g_2g_3$.

(2) *$H \leq A$, $Type(\Gamma(H)) = C_1 *_{\{v_1 \cdot a = v_2 \cdot a \mid a \in A\}} C_2$ and*

$$Type(\Gamma(K)) = \begin{cases} C_l, & K \nleq A; \\ C_l *_{\{v \cdot a = K \cdot a \mid a \in A\}} Cayley(G_j, K), & K \leq A, \end{cases}$$

where $(C_i, v_i) = (Cayley(G_i, H), H \cdot 1)$, for all $i \in \{1, 2\}$, $Lab(C_l, v) = K$, $v \in V(C_l)$ and $1 \leq l \neq j \leq 2$.

Proof. If (1) holds then, by Lemma 5.15, H is conjugate to K in G.

Assume that (2) holds and, without loss of generality, assume that $l = 1$. Thus, by Lemma 5.13 (iii), $Lab(\Gamma(K), v) = Lab(Type(\Gamma(K)), v)$. Therefore, by Lemma 5.5, $Lab(Type(\Gamma(K)), v) = Lab(C_1, v)$. Therefore the subgroup $Lab(C_1, v)$ is conjugate to the subgroup $Lab(\Gamma(K), u_0) = K$.

On the other hand, $Lab(\Gamma(H), v) = Lab(C_1, v)$, by Remark 5.2, because C_2 is redundant w.r.t. v. Therefore the subgroup $Lab(C_1, v)$ is conjugate to the subgroup $Lab(\Gamma(H), v_0) = H$. Thus H and K are conjugate subgroups of G.

Let $\underline{K = g^{-1}Hg}$. Without loss of generality, assume that $g \in G$ is a normal word. Let $g \equiv g_1g_2$, where g_1 is the maximal prefix of the word g such that there is a path p in $\Gamma(H)$ with $\iota(p) = v_0$ and $lab(p) \equiv g_1$, where v_0 is the basepoint of $\Gamma(H)$. Let $v = \tau(p) \in V(\Gamma(H))$. See Figure 7.

If g_2 is the empty word then $g \equiv g_1$, and $Lab(\Gamma(H), v) = g^{-1}Hg = K$.

If $v \in Type(\Gamma(H))$ then, by Lemma 5.13 (iv), we are done.

Assume now that $v \notin V(Type(\Gamma(H))$. Therefore $Type(\Gamma(H)) \neq \Gamma(H)$. Thus, by Definition 5.12, $H \nleq G_i$ $(i \in \{1, 2\})$. Without loss of generality, we can assume that $Type(\Gamma(H)) = \Gamma_m$, where Γ_m is obtained from $\Gamma(H)$ by the iterative removal of the unique sequence $(*)$ of alternating monochromatic components

$$C_0, C_1, \ldots, C_{m-1}.$$

Hence there exists $1 \leq i \leq m-1$ such that $v \in V(C_i)$. Without loss of generality, we can assume that C_i is a X_1-monochromatic component. Let



Figure 7. The closed connected curves represent monochromatic components of different colors. The broken curves denote the rest of the graph.

Γ_i be the graph obtained from $\Gamma(H)$ by the iterative removal of the unique sequence $C_0, \ldots, C_{i-1}$. By Lemma 5.5, we have either $\Gamma(K) = \Gamma_i$ or

$$\Gamma(K) = \Gamma_i *_{\{v \cdot a = S \cdot a \mid a \in A\}} Cayley(G_2, S),$$

where $S = K \cap A$.

In the first case, since each component C_j is redundant in Γ_j $(i \leq j \leq m-1)$, $Type(\Gamma(K))$ is obtained from $\Gamma(K)$ by the iterative erasure of the unique sequence of alternating monochromatic components $C_i, \ldots, C_{m-1}$.

In the second case, the component $D = Cayley(G_2, S)$ of $\Gamma(K)$ is redundant. Therefore $Type(\Gamma(K))$ is obtained from $\Gamma(K)$ by the iterative erasure of the unique sequence of alternating monochromatic components $D, C_i, \ldots, C_{m-1}$. Therefore $K \not\leq G_i$ $(i \in \{1, 2\})$ and

$$Type(\Gamma(K)) = \Gamma_m = Type(\Gamma(H)).$$

Assume now that g_2 is a nonempty word.

We suppose first that $v \in V(Type(\Gamma(H)))$. Let Γ' be the graph obtained from $Type(\Gamma(H))$ by attaching to this graph a "stem" q at the vertex v, such that $lab(q) \equiv g_2$. Thus $\iota(q) = v$ and we let $\tau(q) = v'$, see Figure 8.

Obviously, $Lab(\Gamma', v) = Lab(Type(\Gamma(H)), v)$. By Lemma 5.13,

$$Lab(\Gamma', v) = Lab(Type(\Gamma(H)), v) = Lab(\Gamma(H), v) = g_1^{-1} H g_1.$$

Therefore $Lab(\Gamma', v') = g_2^{-1} Lab(\Gamma', v) g_2 = g^{-1} H g = K$.

Let $q = q_1 \cdots g_k$ be a decomposition of q into maximal monochromatic paths. Let $v_i = \iota(q_i)$, $1 \leq i \leq k$. Thus $v = v_1$.



FIGURE 8. The closed connected curves represent monochromatic components of different colors. The broken curves denote the rest of the graph.

Now we need the following result from [33] (given along with the proof of Claim 2 in [33]).

Claim 2. *The graph* (Γ', v') *can be embedded into a finite precover* (Γ'', v'') *of* G *such that*

$$\Gamma'' = \Big(\big((\Gamma' *_{\{v_1 \cdot a | a \in A\}} D_1\big) *_{\{v_2 \cdot a | a \in A\}} D_2\big) \cdots\Big) *_{\{v_k \cdot a | a \in A\}} D_k,$$

where

- $(D_j, v_j) = Cayley(G_{i_j}, S_j)$ $(1 \leq j \leq k,\ 1 \leq i_j \neq i_{j+1} \leq 2)$*,*
- $S_1 = Lab(C, v) \cap A$*, where* C *is a* X_{i_1} *monochromatic component of* $Type(\Gamma(H))$ *such that* $v \in V(C)$*,*
- $S_{j+1} = Lab(Cayley(G_{i_j}, S_j), v_{j+1}) \cap A$ $(1 \leq j \leq k-1)$*,*
- *the image of* q_j *in* Γ'' *is a path in* D_j*,*
- v'' *is the image of* v' *in* Γ''*.*

Let $\Gamma'_j = \big((\Gamma' *_{\{v_1 \cdot a | a \in A\}} D_1\big) \cdots\big) *_{\{v_j \cdot a | a \in A\}} D_j$, for all $1 \leq j \leq k$. Thus $\Gamma'' = \Gamma'_k$.

By Corollary 5.6, $Lab(\Gamma', v_1) = Lab(\Gamma'_1, v_1)$ and $Lab(\Gamma'_j, v_j) = Lab(\Gamma'_{j-1}, v_j)$.[7] Therefore

$$\begin{aligned} Lab(\Gamma'_j, v_{j+1}) &= (lab(q_j))^{-1} Lab(\Gamma'_j, v_j) lab(q_j) = (lab(q_j))^{-1} Lab(\Gamma'_{j-1}, v_j) lab(q_j) \\ &= (lab(q_1 \cdots q_j))^{-1} Lab(\Gamma'_1, v_1) lab(q_1 \cdots q_j) \\ &= (lab(q_1 \cdots q_j))^{-1} Lab(\Gamma', v_1) lab(q_1 \cdots q_j). \end{aligned}$$

Thus

$$Lab(\Gamma'', v'') = Lab(\Gamma'_k, v'') = g_2^{-1} Lab(\Gamma', v_1) g_2 = (g_1 g_2)^{-1} H (g_1 g_2) = K.$$

Moreover, by the construction, Γ'' is a precover of G which has no redundant components w.r.t. v''. Hence, by Lemma 5.5, either $\Gamma(K) = \Gamma''$ or

$$\Gamma(K) = \Gamma'' *_{\{v'' \cdot a = S \cdot a \mid a \in A\}} Cayley(G_l, S),$$

where $S = K \cap A$ and $l = i_{k-1}$.

By the construction of Γ'' (see the proof of Claim 2), $D_k, \ldots, D_1$ is the unique sequence of redundant components in Γ'' which satisfies the conditions (1)-(3) from the description of sequence $(*)$ (see the proof of Lemma 5.11). Therefore, in the first case, it should be erased from $\Gamma(K)$ along the construction of $Type(\Gamma(K))$.

In the second case, the component $D = Cayley(G_2, S)$ of $\Gamma(K)$ is redundant. Therefore the sequence $D, D_k, \ldots, D_1$ should be erased from $\Gamma(K)$ along the construction of $Type(\Gamma(K))$. Therefore, by Definition 5.12, $K \neq G_i$ $(i \in \{1,2\})$. Moreover, if $Type(\Gamma(H))$ has no redundant components, that is $H \not\leq A$ then $Type(\Gamma(K)) = Type(\Gamma(H))$ if $Type(\Gamma(H))$. Otherwise

$$Type(\Gamma(H)) = \Gamma(H) = C_1 *_{\{v_1 \cdot a = v_2 \cdot a \mid a \in A\}} C_2,$$

where $(C_i, v_i) = (Cayley(G_i, H), H \cdot 1)$, for all $i \in \{1,2\}$. Without loss of generality assume that $v \in V(C_1)$. Hence $Type(\Gamma(K)) = C_1$.

If $v \notin V(Type(\Gamma(H)))$ then $H \not\leq G_i$ $(i \in \{1,2\})$. We take Γ' to be the graph obtained by gluing a stem q labelled by g_2 at $v \in V(C_i)$ to the graph Γ_i, which is obtained from $\Gamma(H)$ by the iterative removal of the sequence $C_0, \ldots, C_{i-1}$ of redundant components in $\Gamma(H)$. Combining the proofs of two previous cases, namely $v \notin V(Type(\Gamma(H)))$, $g_2 =_G 1$ and $v \in V(Type(\Gamma(H)))$, $g_2 \neq_G 1$, we conclude that $K \not\leq G_i$ $(i \in \{1,2\})$, and $Type(\Gamma(K)) = Type(\Gamma(H))$.

◇

---

[7] By abuse of notation, we identify the vertices $v_j \in V(\Gamma')$ with their images in the graphs Γ'_j $(1 \leq j \leq k)$.

Corollary 5.17. *Let $h_1, \ldots, h_s, k_1, \ldots, k_t \in G$. Then there is an algorithm, which decides whether or not the subgroups*

$$H = \langle h_1, \ldots, h_s \rangle \text{ and } K = \langle k_1, \ldots, k_t \rangle \quad (i \in \{1, 2\})$$

are conjugate in G.

Proof. First we construct the graphs $(\Gamma(H), v_0)$ and $(\Gamma(K), u_0)$, using the generalized Stallings' algorithm. Then we compute $Type(\Gamma(H))$ and $Type(\Gamma(K))$ according to the definition of $Type$. Now we verify if any of the conditions from Theorem 5.16 are satisfied.

Note that the verification of $\Delta_1 = \Delta_2$ actually means to check whether or not Γ_1 and Γ_2 are isomorphic. This can be done by fixing a vertex $v \in V(\Delta_1)$ and comparing for each vertex $w \in V(\Delta_2)$ the pointed graphs (Δ_1, v) and (Δ_2, w), because by Remark 3.1, such an isomorphism if it exists is unique.

Since morphisms of well-labelled graphs preserves endpoints and labels, we can specify the above verification by fixing a bichromatic vertex $v \in VB(\Delta_1)$ and comparing the pointed graphs (Δ_1, v) and (Δ, w), for each bichromatic vertex $w \in VB(\Gamma_2)$.

◇

Example 5.18. The subgroups H_1 and H_2 from Example A.2 (see Figures 12 and 13) are not conjugate to each other, because their $Type$ graphs are not isomorphic. Indeed, $Type(\Gamma(H_i)) = \Gamma(H_i)$ for $i \in \{1, 2\}$, but $|V(\Gamma(H_1))| \neq |V(\Gamma(H_2))|$. Hence these graphs can not be isomorphic. ◇

Complexity. Let m be the sum of the lengths of the words $h_1, \ldots h_s$, and let l be the sum of the lengths of the words $k_1, \ldots, k_t$. By Theorem 4.1 (5), the complexity of the construction of the graphs $\Gamma(H)$ and $\Gamma(K)$ is $O(m^2)$ and $O(l^2)$, respectively.

The detecting of monochromatic components in the constructed graphs takes $O(|E(\Gamma(H))|)$ and $O(|E(\Gamma(K))|)$, that is $O(m)$ and $O(l)$, respectively. Since all the essential information about A, G_1 and G_2 is given and it is not a part of the input, verifications concerning a particular monochromatic component of $\Gamma(H)$ or of $\Gamma(K)$ takes $O(1)$. Therefore, the complexity of the construction of $Type(\Gamma(H))$ from $\Gamma(H)$ is $O(|E(\Gamma(H))|)$, that is $O(m)$. Similarly, the complexity of the construction of $Type(\Gamma(K))$ from $\Gamma(K)$ is $O(|E(\Gamma(K))|)$, that is $O(l)$.

Now we are ready to verify an isomorphism of the obtained type graphs. We can start by comparing the sizes of $V(\Gamma_1)$ and $V(\Gamma_2)$ and of $E(\Gamma_1)$ and $E(\Gamma_2)$. If $|V(\Gamma_1)| = |V(\Gamma_2)|$ and $|E(\Gamma_1)| = |E(\Gamma_2)|$ then we continue. Otherwise the graphs are not isomorphic.

Let $\Gamma_1 = Type(\Gamma(H))$ and $\Gamma_2 = Type(\Gamma(K))$. Let $v \in VB(\Gamma_1)$ and $w \in VB(\Gamma_2)$. Thus, by Definition 5.4, (Γ_1, v) and (Γ_2, w) are finite reduced precovers of G.

Theorem 4.1 (2) implies that the finite reduced precovers (Γ_1, v) and (Γ_2, w) are isomorphic if and only if they are isomorphic via the morphism μ of well-labelled pointed graphs, defined in the proof of Lemma 4.6 in [11]. That is to check the isomorphism between (Γ_1, v) and (Γ_2, w), we simply have to check if μ is defined. Recall that $\mu : (\Gamma_1, v) \to (\Gamma_2, w)$ satisfies

$$\mu(\vartheta) = w \cdot x \ \big(\forall\, \vartheta = v \cdot x \in V(\Gamma_1)\big) \ \text{ and } \ \mu(e) = (\mu(\iota(e)), lab(e)) \ \big(\forall\, e \in E(\Gamma)\big).$$

Thus for all $\vartheta = v \cdot x \in V(\Gamma_1)$ we have to check if $Star(v \cdot x, \Gamma_1) = Star(w \cdot x, \Gamma_2)$, where the *star* of the vertex σ (see [46]) in the graph Δ is the set

$$Star(\sigma, \Delta) = \{e \in E(\Delta) \mid \iota(e) = \sigma\}.$$

This procedure takes time proportional to $|E(\Gamma_1)|$, that is proportional to m. Since in the worst case we have to repeat the above procedure for all pointed graphs (Γ_2, ω), where $\omega \in VB(\Gamma_2)$, the verification of an isomorphism between the graphs Γ_1 and Γ_2 takes $O\big(|VB(\Gamma_2)| \cdot |E(\Gamma_1)|\big)$.

Since $|VB(\Gamma_2)| \leq |V(\Gamma_2)|$ and, by Theorem 4.1 (5), $|V(\Gamma_2)|$ is proportional to l and $|E(\Gamma_1)|$ is proportional to m, the complexity of the algorithm given along with the proof of Corollary 5.17 is

$$O\big(m^2 + l^2 + ml\big) \ = \ O\big((m + l)^2\big).$$

Thus the above algorithm is quadratic in the size of the input.

Note that if the subgroups H and K are given by the graphs $\Gamma(H)$ and $\Gamma(K)$, the complexity of the algorithm that decides whether or not the subgroup H and K are conjugate in G is

$$O\big(|E(\Gamma(H))|^2 + |E(\Gamma(K))|^2 + |VB(\Gamma_2)| \cdot |E(\Gamma_1)|\big).$$

Note that, since our graphs are connected, $|V(\Gamma_2)| \leq |E(\Gamma_2)|$. Thus $|VB(\Gamma_2)| \leq |V(\Gamma_2)| \leq |E(\Gamma_2)| \leq |E(\Gamma(K))|$. Since $|E(\Gamma_1)| \leq |E(\Gamma(H))|$, the complexity is

$$O\big(|E(\Gamma(H))|^2 + |E(\Gamma(K))|^2 + |E(\Gamma(H))| \cdot |E(\Gamma(K))|\big).$$

That is it is quadratic in the size of the input $O\Big(\big(|E(\Gamma(H))| + |E(\Gamma(K))|\big)^2\Big)$.

5.3. **More Conjugacy Results.**

Theorem 5.19. *Let H and K be finitely generated subgroup of an amalgam of finite groups $G = G_1 *_A G_2$.*
Then there exists $g \in G$ such that $gKg^{-1} \leq H$ if and only if one of the following holds.

(1) *If* $K \nleq A$ *then there exists a morphism of well-labelled graphs*

$$\pi : Type(\Gamma(K)) \rightarrow Type(\Gamma(H)).$$

(2) *If* $K \leq A$ *and* $Type(\Gamma(K)) = C_1 *_{\{v_1 \cdot a = v_2 \cdot a \mid a \in A\}} C_2$*, where* $(C_i, v_i) = (Cayley(G_i, K), K \cdot 1)$ $(i \in \{1,2\})$*, then there exists a morphism of well-labelled graphs* $\pi : C_l \rightarrow Type(\Gamma(H))$*, for some* $l \in \{1,2\}$.

Proof. We begin with the following claim which allows to assume that $\Gamma(H) = Type(\Gamma(H))$.

Claim 3. *There exist* $v \in V(\Gamma(H))$ *such that* $Type(\Gamma(L)) = \Gamma(L)$*, where* $L = Lab(Type(\Gamma(H)), v)$.

Proof of the Claim. By Definition 5.12, either $Type(\Gamma(H)) = \Gamma(H)$ or $Type(\Gamma(H)) = \Gamma_m$.

If $\Gamma_m = C_m$ then we take $v \in VB_{\Gamma_{m-1}}(C_{m-1}) = VB_{\Gamma_{m-1}}(C_m)$. Thus $L = Lab(C_m, v) \leq G_i$ $(i \in \{1,2\})$, and, by the proof of Lemma 5.11, $L \nleq A$. Hence, by Lemma 5.7, $\Gamma(L) = C_m$. Thus, by Definition 5.12, $\Gamma(L) = Type(\Gamma(L))$.

Assume now that Γ_m has at least two distinct monochromatic components. Let $v \in VB(\Gamma_m)$. By Lemma 5.13(i), Γ_m is a finite nonempty precover of G. Thus, since Γ_m has no redundant components and $v \in VB(\Gamma_m)$, we conclude that Γ_m is a finite reduced precover. That is $(\Gamma_m, v) = (\Gamma(L), u_0)$, where u_0 is the basepoint of $\Gamma(L)$. Therefore $Type(\Gamma(L)) = \Gamma(L)$.

◇

Let $y \in G$ such that $v = v_0 \cdot y$. Therefore $L = y^{-1}Hy$. Thus $y^{-1}gKg^{-1}y \leq L$. Hence, without loss of generality, we can assume that $\Gamma(H) = Type(\Gamma(H))$.

Assume first that there exists $g \in G$ such that $gKg^{-1} \leq H$. By Lemma 3.2, there exists a morphism $\varphi : \Gamma(gKg^{-1}) \rightarrow \Gamma(H)$. Let φ' be the restriction of φ to $Type(\Gamma(gKg^{-1}))$, that is $\varphi' : Type(\Gamma(gKg^{-1})) \rightarrow Type(\Gamma(H))$.

Let $K \nleq A$. Thus either $gKg^{-1} \nleq A$ or $gKg^{-1} \leq A$. Hence, by Theorem 5.16, either $Type(\Gamma(K)) = Type(\Gamma(gKg^{-1}))$ or,

$$Type(\Gamma(gKg^{-1})) = C_1 *_{\{v_1 \cdot a = v_2 \cdot a \mid a \in A\}} C_2, \quad Type(\Gamma(K)) = C_l \ (l \in \{1,2\}),$$

where $(C_i, v_i) = (Cayley(G_i, gKg^{-1}), gKg^{-1} \cdot 1)$, $i \in \{1,2\}$. In the first case, we take $\pi = \varphi'$. In the second one, there exists an embedding $\phi : Type(\Gamma(K)) \rightarrow Type(\Gamma(gKg^{-1}))$. Therefore $\pi = \varphi' \circ \phi$ gives the desired morphism.

Assume now that $K \leq A$. Thus, by Definition 5.12 and Lemma 5.7 (iii), $Type(\Gamma(K)) = C_1 *_{\{v_1 \cdot a = v_2 \cdot a \mid a \in A\}} C_2$, where $(C_i, v_i) = (Cayley(G_i, K), K \cdot 1)$ $(i \in \{1,2\})$.

If $gKg^{-1} \nleq A$ then, by Theorem 5.16, $Type(\Gamma(gKg^{-1})) = C_l$, for some $l \in \{1, 2\}$. Thus $\pi = \varphi'$ produces the desired morphism $\pi : C_l \to Type(\Gamma(H))$. If $gKg^{-1} \leq A$ then, by Theorem 5.16,

$$Type(\Gamma(gKg^{-1})) = C_l *_{\{v \cdot a = (gKg^{-1}) \cdot a \mid a \in A\}} Cayley(G_j, gKg^{-1}),$$

where $1 \leq l \neq j \leq 2$. Thus there exists an embedding $\phi : C_l \to Type(\Gamma(gKg^{-1}))$. Therefore $\pi = \varphi' \circ \phi : C_l \to Type(\Gamma(H))$ gives the desired morphism.

Suppose now that $K \nleq A$ and the morphism $\pi : Type(\Gamma(K)) \to Type(\Gamma(H))$ exists. Let p be a path in $\Gamma(K)$ with $\iota(p) = u_0$, where u_0 is the basepoint of the graph $\Gamma(K)$, such that $\tau(p) \in V(Type(\Gamma(K)))$. Let $u = \tau(p)$, $lab(p) \equiv f$ and let $\vartheta = \pi(u) \in V(Type(\Gamma(H)))$.

Since $Type(\Gamma(H)) \subseteq \Gamma(H)$, we have $\vartheta \in V(\Gamma(H))$. By Lemma 4.10, there exists a normal path q in $\Gamma(H)$ with $\iota(q) = v_0$ (the basepoint of $\Gamma(H)$) and $\tau(q) = \vartheta$. Let $lab(q) \equiv c$. By Lemma 5.13 (iii), $Lab(Type(\Gamma(K)), u) = Lab(\Gamma(K), u) = f^{-1}Kf$ and $Lab(Type(\Gamma(H)), \vartheta) = Lab(\Gamma(H), \vartheta) = c^{-1}Hc$.

Since π can also be considered as a morphism of pointed graphs

$$\pi : (Type(\Gamma(K)), u) \to (Type(\Gamma(H)), \vartheta),$$

by Lemma 3.2, we have $Lab(Type(\Gamma(K)), u) \leq Lab(Type(\Gamma(H)), \vartheta)$. Thus $f^{-1}Kf \leq c^{-1}Hc$. Therefore $g = c \cdot f^{-1}$ and $gKg^{-1} \leq H$, as required.

Let $K \leq A$. Thus, by Lemma 5.7 (iii), $\Gamma(K) = C_1 *_{\{v_1 \cdot a = v_2 \cdot a \mid a \in A\}} C_2$, where $(C_i, v_i) = (Cayley(G_i, K), K \cdot 1)$ $(i \in \{1, 2\})$ and the basepoint u_0 of $\Gamma(K)$ is the image of $K \cdot 1$. Thus $Lab(C_1, u_0 \cdot a) = Lab(C_2, u_0 \cdot a) = Lab(\Gamma(K), u_0 \cdot a)$, for all $a \in A$.

Assume that there is $l \in \{1, 2\}$ such that the morphism $\pi : C_l \to Type(\Gamma(H))$ exists. Let p be a path in $\Gamma(K)$ with $\iota(p) = u_0$, where u_0 is the basepoint of the graph $\Gamma(K)$, such that $\tau(p) \in V(C_l)$. Let $u = \tau(p)$, $lab(p) \equiv f$ and let $\vartheta = \pi(u) \in V(Type(\Gamma(H)))$.

Since $Type(\Gamma(H)) \subseteq \Gamma(H)$, we have $\vartheta \in V(\Gamma(H))$. By Lemma 4.10, there exists a normal path q in $\Gamma(H)$ with $\iota(q) = v_0$ (the basepoint of $\Gamma(H)$) and $\tau(q) = \vartheta$. Let $lab(q) \equiv c$. Thus $Lab(C_l, u) = f^{-1}Lab(C_l, u_0)f = f^{-1}Lab(\Gamma(K), u_0)f = f^{-1}Kf$ and $Lab(Type(\Gamma(H)), \vartheta) = Lab(\Gamma(H), \vartheta) = c^{-1}Hc$.

Since π can also be considered as a morphism of pointed graphs

$$\pi : (C_l, u) \to (Type(\Gamma(H)), \vartheta),$$

by Lemma 3.2, we have $Lab(C_l, u) \leq Lab(Type(\Gamma(H)), \vartheta)$. Thus $f^{-1}Kf \leq c^{-1}Hc$. Therefore $g = c \cdot f^{-1}$ and $gKg^{-1} \leq H$, as required.

◇

Corollary 5.20. *Let* $h_1, \ldots, h_s, k_1, \ldots, k_t \in G$. *Then there exists an algorithm which decides whether or not there exists* $g \in G$ *such that* $gKg^{-1} \leq H$, *where*

$$H = \langle h_1, \ldots, h_s \rangle \text{ and } \; K = \langle k_1, \ldots, k_t \rangle.$$

Moreover, the algorithm produces one such g if it exists.

Proof. First we construct the graphs $(\Gamma(K), u_0)$ and $(\Gamma(H), v_0)$, using the generalized Stallings' folding algorithm. Then we construct $Type(\Gamma(K))$ and $Type(\Gamma(H))$, according to Definition 5.12.

If $K \not\leq A$ then we proceed as follows. Let $u \in V(Type(\Gamma(K)))$. For each vertex $v \in V(Type(\Gamma(H)))$ we iteratively check if there exists a morphism $\pi : (Type(\Gamma(K)), u) \rightarrow (Type(\Gamma(H)), v)$. If no such morphism can be found then K is not conjugate to any subgroup of H, by Theorem 5.19. Otherwise, by the proof of Theorem 5.19, $gKg^{-1} \leq H$, where $g = c \cdot f^{-1}$ and $v = v_0 \cdot f$ and $u = u_0 \cdot c$.

Assume now that $K \leq A$. Thus $Type(\Gamma(K)) = C_1 *_{\{v_1 \cdot a = v_2 \cdot a \mid a \in A\}} C_2$, where $(C_i, v_i) = (Cayley(G_i, K), K \cdot 1)$ $(i \in \{1, 2\})$.

For each $i \in \{1, 2\}$, let $u_i \in V(C_i)$. For each vertex $v \in V(Type(\Gamma(H)))$ we iteratively check if there exists a morphism $\pi : (C_i, u_i) \rightarrow (Type(\Gamma(H)), v)$. If no such morphism can be found then K is not conjugate to any subgroup of H, by Theorem 5.19. Otherwise, by the proof of Theorem 5.19, $gKg^{-1} \leq H$, where $g = c \cdot f^{-1}$ and $v = v_0 \cdot f$ and $u = u_0 \cdot c$.

◇

Complexity. Similarly to the complexity analysis of the algorithm presented along with the proof of Corollary 5.20, the complexity of the above algorithm is $O((m + l)^2)$, where m is the sum of the lengths of the words $h_1, \ldots h_s$, and l is the sum of the lengths of the words $k_1, \ldots, k_t$. Similarly, when the subgroup H and K are given by the graphs $\Gamma(H)$ and $\Gamma(K)$, the complexity is $O\big((|E(\Gamma(H))| + |E(\Gamma(K))|)^2\big)$.

Corollary 5.21 (The Conjugacy Problem)**.** *The conjugacy problem is solvable in amalgams of finite groups.*

Namely, let $G = G_1 *_A G_2$ be an amalgam of finite groups. Given elements $k, h \in G$ one can decide whether exists $g \in G$ such that $gkg^{-1} =_G h$.

Proof. Let $K = \langle k \rangle$ and $H = \langle h \rangle$. We apply to K and H the algorithm described along with the proof of Corollary 5.20. If there is no $g \in G$ such that $gKg^{-1} \leq H$ then the elements k and h are not conjugate in G.

Otherwise, let $g \in G$ such that $gKg^{-1} \leq H$. We have to check whether $gkg^{-1} =_G h$. To this end we rewrite the element $gkg^{-1}h^{-1}$ as a normal word. If the resulting word is not empty then, by the Torsion Theorem (IV.2.7, [27]), $gkg^{-1}h^{-1} \neq_G 1$, that is $gkg^{-1} \neq_G h$. Otherwise, $gkg^{-1} =_G h$. ◇

6. The Normality Problem

The current section is devoted to the solution of the *normality problem*, which asks to know if a subgroup H of a group G is normal in G, for finitely generated subgroups of amalgams of finite groups.

The quadratic time algorithm is presented in Corollary 6.6. It is based on Theorem 6.2 and Lemma 6.3, which give a connection between the normality of a subgroup H of an amalgam of finite groups $G = G_1 *_A G_2$ and its subgroup graph $\Gamma(H)$ constructed by the generalized Stallings' algorithm. The complexity analysis of the algorithm is given at the end of this section.

We start by presenting the following technical result from [32], which is essential for the proof of Theorem 6.2.

Lemma 6.1 (Lemma 6.10 and Remark 6.11 in [32])**.** *Let* $G = G_1 *_A G_2$ *be an amalgam of finite groups. Let* (Γ, v) *be a finite precover of* G *such that* $Lab(\Gamma, v) =_G H \neq \{1\}$.

Let $w \in H$ *be a normal word. Then* w *labels a path in* Γ *closed at* v *if one of the following holds*

- $l(w) > 1$,
- $l(w) = 1$ *and* $w \in G_i \setminus A$ $(i \in \{1, 2\})$,
- $l(w) = 1$, $w \in G_i \cap A$ *and,* $v \in VB(\Gamma)$ *or* $v \in VM_i(\Gamma)$.

Otherwise, if $w \in G_i \cap A$ *and* $v \in VM_j(\Gamma)$ $(1 \leq i \neq j \leq 2)$, *then there exists a path in* Γ *closed at* v *and labelled with* w' *such that* $w' \in G_j \cap A$, $w =_G w'$.

Theorem 6.2. *Let* $H \leq G$ *be a nontrivial subgroup of* G *such that* $H \not\leq G_i$ *for all* $i \in \{1, 2\}$. *Then* H *is normal in* G *if and only if the following holds.*

(i) *The graph* $\Gamma(H)$ *is* $X^{\pm}$*-saturated.*
(ii) *For all vertices* $v, u \in V(\Gamma(H))$, *the graphs* $(\Gamma(H), v)$ *and* $(\Gamma(H), u)$ *are isomorphic.*

Proof. Suppose first that conditions (i) and (ii) are satisfied.

Let g be an element of G. Since $\Gamma(H)$ is $X^{\pm}$-saturated, there exists a path p in $\Gamma(H)$ such that $\iota(p) = v_0$ and $lab(p) \equiv g$. Let $v = \tau(p)$. Condition (ii) implies $Lab(\Gamma(H), v_0) = Lab(\Gamma(H), v)$. Thus $H = g^{-1}Hg$, for all $g \in G$. Hence $H \trianglelefteq G$.

Assume now that $\{1\} \neq H \trianglelefteq G$. Then $\Gamma(H)$ is $X^{\pm}$-saturated. Otherwise, without loss of generality, we can assume that there exists $v \in VM_1(\Gamma(H))$. Let C be the X_1-monochromatic component of $\Gamma(H)$ such that $v \in V(C)$.

Let q be the approach path in $\Gamma(H)$ from v_0 to v with $lab(q) \equiv g$. Thus

$$Lab(\Gamma(H), v) = gLab(\Gamma(H), v_0)g^{-1} = gHg^{-1} = H.$$

Since $(\Gamma(H), v)$ is a precover of G, each normal element of H, whose syllable length is greater than 1, labels a normal path in $(\Gamma(H), v)$ closed at v, by Lemma 6.1.

Let $h \in H$ has the normal decomposition $(h_1, \ldots, h_k)$. Thus $k > 1$, since $H \nleq G_i$ $(i \in \{1, 2\})$. Let p be a normal path in $\Gamma(H)$ such that

$$\iota(p) = \tau(p) = v, \ p = p_1 \cdots p_k, \ \text{where } lab(p_l) \equiv h_l, \ 1 \leq l \leq k.$$

Thus $h_1, h_k \in G_1 \setminus A$, because $v \in VM_1(\Gamma(H))$. Hence $p_k p_1$ is a path in C from $\iota(p_k)$ to $\tau(p_1)$, and we have $lab(p_k p_1) \equiv h_k h_1 \in G_1$.

If $h_k h_1 \notin A$ then the decomposition $(h_2, \ldots, (h_k h_1))$ is normal. Moreover, $h_1^{-1} h h_1 =_G h_2 \cdots (h_k h_1) \in H$, because $H \trianglelefteq G$. Therefore, by Lemma 6.1, there exists a normal path in $\Gamma(H)$ closed at v and labelled with $h_2 \cdots h_{k-2}(h_k h_1)$. However, this is impossible because $v \in VM_1(\Gamma(H))$ and $h_2 \in G_2 \setminus A$. We get a contradiction.

If $h_k h_1 \in G_1 \cap A$, we take $b =_G h_k h_1$ such that $b \in G_2 \cap A$. Thus the decomposition $(h_2, \ldots, (h_{k-1} b))$ is normal, since $h_{k-1} b \in G_2 \setminus A$. We get a contradiction in the similar way.

Therefore the graph $\Gamma(H)$ is $X^{\pm}$-saturated. Moreover,

$$Lab(\Gamma(H), v) = gLab(\Gamma(H), v_0) g^{-1} = gHg^{-1} = H,$$

where $g \equiv lab(q)$ and q is an approach path in $\Gamma(H)$ from v_0 to v.

Thus, by Lemma 4.6, $(\Gamma(H), v)$ is isomorphic to $(Cayley(G, H), H \cdot 1)$, for all $v \in V(\Gamma(H))$. Therefore the graphs $(\Gamma(H), v)$ and $(\Gamma(H), u)$ are isomorphic, for all vertices $v, u \in V(\Gamma(H))$.

◇

Lemma 6.3. *Let H be a nontrivial subgroup of G such that $H \leq G_i$ $(i \in \{1, 2\})$. The following holds.*

- (i) *If $H \trianglelefteq G$ then $H \leq A$.*
- (ii) *If $H \leq A$ then $H \trianglelefteq G$ if and only if each monochromatic component C of $\Gamma(H)$ is a regular graph, that is (C, v) is isomorphic to (C, u), for all $v, u \in V(C)$.*

Proof. To prove (i) suppose that there exists $h \in H \setminus A$. Let $g \in G_j \setminus A$, where $1 \leq i \neq j \leq 2$. Therefore ghg^{-1} is a normal word of syllable length 3. Hence $ghg^{-1} \notin H$. This contradicts with the assumption that $H \trianglelefteq G$.

Since $H \leq A$, $\Gamma(H) = Cayley(G_1, H) *_{\{Ha \mid a \in A\}} Cayley(G_2, H)$, by Lemma 5.7 (iii). Since $H \leq A$, $H \trianglelefteq G$ if and only if $H \trianglelefteq G_i$ $(i \in \{1, 2\})$. Therefore $H \trianglelefteq G$ if and only if the graphs $Cayley(G_1, H)$ and $Cayley(G_2, H)$ are regular (see 2.2.7 in [47]). ◇

Recall the following result from [33].

Theorem 6.4 (Theorem 7.1 in [33]). *Let H be a finitely generated subgroup of an amalgam of finite groups $G = G_1 * G_2$.*

Then $[G : H] < \infty$ if and only if $\Gamma(H)$ is $X^{\pm}$-saturated.

Remark 6.5. *By Theorem 6.4, $H \trianglelefteq G$ implies $[G : H] < \infty$.*

Corollary 6.6. *Let $h_1, \dots h_k \in G$. Then there exists an algorithm which decides whether or not $H = \langle h_1, \dots, h_k \rangle$ is a normal subgroup (of finite index) in G.*

Proof. We first construct the graph $\Gamma(H)$ using the generalized Stallings' algorithm.

If the number of monochromatic components of $\Gamma(H)$ is equal to 1 then, by Lemma 5.7 (ii), $H \leq G_i$ and $H \cap A = \{1\}$ ($i \in \{1, 2\}$). Hence, by Lemma 6.3 (i), H is not normal in G.

If the number of distinct monochromatic components of $\Gamma(H)$ is equal 2 and

$$(\Gamma(H), v_0) = (Cayley(G_1, H) *_{\{Ha \mid a \in A\}} Cayley(G_2, H), \vartheta),$$

where ϑ is the image of $H \cdot 1$ in the amalgam graph, then $H \leq A$, by Lemma 5.7 (iii). Thus, by Lemma 6.3 (ii), H is normal in G if and only if $Cayley(G_1, H)$ and $Cayley(G_2, H)$ are regular graphs, that is if and only if $(Cayley(G_i, H), v)$ is isomorphic to $(Cayley(G_i, H), H \cdot 1)$, for all $v \in V(Cayley(G_i, H))$, $i \in \{1, 2\}$. Since an isomorphism of pointed labelled graphs is unique, by Remark 3.1, we are done.

Assume now that $H \not\leq G_i$, Thus we verify if $\Gamma(H)$ is $X^{\pm}$-saturated. If it is not then, by Theorem 6.2, H is not normal in G. Otherwise, if $\Gamma(H)$ is $X^{\pm}$-saturated, then, by Theorem 6.2, H is normal in G if and only if the graphs $(\Gamma(H), v_0)$ and $(\Gamma(H), v)$ are isomorphic, for each vertex $v \in V(\Gamma(H))$. Since an isomorphism of pointed labelled graphs is unique, by Remark 3.1, we are done.

◇

Example 6.7. Let H_1 and H_2 be subgroups from Example A.2. One can easily verify from Figures 12 and 13 that H_1 is not normal in G, because $\Gamma(H_1)$ is not $\{x, y\}^{\pm}$-saturated, while $H_2 \trianglelefteq G$. ◇

Complexity. By Theorem 4.1 (5), the complexity of the construction of $\Gamma(H)$ is $O(m^2)$, where m is the sum of lengths of the given subgroup generators.

The detecting of monochromatic components in the constructed graph takes $O(|E(\Gamma(H))|)$, that is $O(m)$, by Theorem 4.1 (5).

Since all the essential information about the amalgam $G = G_1 *_A G_2$, A, G_1 and G_2 is given and it is not a part of the input, the verifications concerning monochromatic components of $\Gamma(H)$ takes $O(1)$. Therefore, to check from $\Gamma(H)$ whether $H \leq G_i$, or $H \leq A$ and the monochromatic components of $\Gamma(H)$ are regular graphs, takes $O(1)$.

To verify that all the vertices of $\Gamma(H)$ are bichromatic takes $O(|E(\Gamma(H))|)$. The verification of an isomorphism of the graphs $(\Gamma(H), v_0)$ and $(\Gamma(H), v)$, for all $v \in V(\Gamma)$, takes time proportional to $|V(\Gamma(H))| \cdot |E(\Gamma(H))|$ (see the complexity analysis of the conjugacy problem). Since, by the Theorem 4.1 (5), $|V(\Gamma(H))|$ and $|E(\Gamma(H))|$ are proportional to m, the complexity of the above algorithm is $O(m^2)$.

If the subgroup H is given by the graph $(\Gamma(H), v_0)$ and not by a finite set of subgroup generators, then the complexity is equal to $|V(\Gamma(H))| \cdot |E(\Gamma(H))|$. Thus in both cases the algorithm is quadratic in the size of the input.

7. Intersection Properties

In this section we study properties of intersections of finitely generated subgroups of amalgams of finite groups such as the *Howson property* (the intersection of two finitely generated subgroups is finitely generated), *malnormality* and *almost malnormality*. The corresponding algorithmic problems and their solutions are presented.

Definition 7.1. *Let H be a subgroup of a group G.*

We say that H is a malnormal subgroup of G if and only if

$$gHg^{-1} \cap H = \{1\}, \ \forall g \in G \setminus H.$$

H is almost malnormal if for all $g \in G \setminus H$, the subgroup $H \cap gHg^{-1}$ is finite.

Obviously, $\{1\}$ and G are malnormal subgroups of G. If G is Abelian, then $\{1\}$ and G are the only malnormal subgroups of G. The most natural example of a malnormal subgroup is K inside any free product $K * L$.

Malnormal subgroups of hyperbolic groups have recently become the object of intensive studies (see, for instance, [10, 11, 12, 14]). Thus malnormality plays an important role in the Combination Theorem for hyperbolic groups [1, 13, 23]. For importance of almost malnormality see, for example, [39, 50].

As is well known [6], in general, malnormality is undecidable in hyperbolic groups. However, the results presented in the current section show that malnormality is *decidable* for finitely generated subgroups in the class of amalgams of finite groups. Below we present a polynomial time algorithm (Corollary 7.12) that solves the *malnormality problem*, which asks to decide whether or not a subgroup H of the group G is malnormal in G. The complexity analysis of the presented algorithm is given at the end of this section.

Product graphs (Definition 7.2) are used to compute intersections of subgroups via their subgroup graphs. We start by studding products of precovers. As an immediate consequence, a solution of *the intersection problem*, which asks to find effectively the intersection of two subgroups, is obtained (Corollary 7.7). This allows one to conclude (Corollary 7.9) that amalgams of finite groups possess *Howson property*, which is known to be true (see, for instance [10, 44]).

Then we characterize malnormality of a finitely generated subgroup H of an amalgam of finite groups by the properties of the product graph $\Gamma(H) \times \Gamma(H)$ (Theorem 7.11). This provides the solution of the malnormality problem (Corollary 7.12). These results are naturally extended to detect *almost malnormality* of finitely generated subgroups of amalgams of finite groups (Theorem 7.14 and Corollary 7.15).

Product Graphs.

Definition 7.2. *Let Γ and Δ be two labelled with $X_1^{\pm} \cup X_2^{\pm}$ graphs. The* *product graph* *$\Gamma \times \Delta$ is the graph defined as follows.*

(1) $V(\Gamma \times \Delta) = V(\Gamma) \times V(\Delta)$.

(2) *for each pair of vertices (v_1, u_1), $(v_2, u_2) \in V(\Gamma \times \Delta)$ (so that $v_1, v_2 \in V(\Gamma)$ and $u_1, u_2 \in V(\Delta)$) and the letter $x \in X$ there exists an edge $e \in E(\Gamma \times \Delta)$ with*

$$\iota(e) = (v_1, u_1), \ \tau(e) = (v_2, u_2) \text{ and } lab(e) \equiv x$$

if and only if there exist edges $e_1 \in E(\Gamma)$ and $e_2 \in E(\Delta)$ such that

$$\iota(e_1) = v_1, \ \tau(e_1) = v_2 \text{ and } lab(e_1) \equiv x$$

and

$$\iota(e_2) = u_1, \ \tau(e_2) = u_2 \text{ and } lab(e_2) \equiv x$$

Along this section we consider $G = G_1 *_A G_2$ to be an amalgam of finite groups G_1 and G_2.

Lemma 7.3. *Let Γ and Δ be finite precovers of $G = G_1 *_A G_2$.*

Then nonempty connected components of $\Gamma \times \Delta$ are finite precovers of G.

Proof. Since Γ and Δ are finite graphs the product graph is finite, by Definition 7.2. Thus each of its connected components is finite as well.

Let Φ be a nonempty connected component Φ of $\Gamma \times \Delta$, that is $E(\Phi) \neq \emptyset$.

Let p be a path in Φ such that $lab(p) \equiv r$, where $r =_G 1$. Let $(v, u) = \iota(p)$ and $(v', u') = \tau(p)$ then there exist paths p_1 in Γ and p_2 in Δ such that

$$\iota(p_1) = v, \ \tau(p_1) = v' \text{ and } lab(p_1) \equiv r$$

and

$$\iota(p_2) = u, \ \tau(p_2) = u' \text{ and } lab(p_2) \equiv r.$$

Since Γ and Δ are G-based graphs, we have $v = \iota(p_1) = \tau(p_1) = v'$ and $u = \iota(p_2) = \tau(p_2) = u'$. Thus $(v, u) = (v', u')$. Hence p is a closed path in Φ. That is Φ is G-based.

Finally, we have to show that each X_i-monochromatic component of Φ is a cover of G_i ($i \in \{1, 2\}$). By Lemma 4.6, a X_i-monochromatic component is a cover of G_i if and only if it is $X_i^{\pm}$-saturated and G_i-based ($i \in \{1, 2\}$).

Let C be a X_i-monochromatic component of Φ. Since the graph Φ is G-based, it is, in particular, G_i-based. Hence C is G_i-based.

C is $X_i^{\pm}$-saturated. Indeed, let $(v, u) \in V(C)$. Thus either $(v, u) \in VM_i(C)$ ($i \in \{1, 2\}$) or $(v, u) \in VB(C)$. Definition 7.2 implies that in the first case at least one of the vertices v or u is X_i-monochromatic in Γ and Δ, respectively, and the other one is bichromatic or X_i-monochromatic. If $(v, u) \in VB(C)$ then $v \in VB(\Gamma)$ and $u \in VB(\Delta)$.

Since Γ and Δ are precovers, their bichromatic vertices are $X^{\pm}$-saturated and their X_i-monochromatic vertices are $X_i^{\pm}$-saturated ($i \in \{1, 2\}$).

Therefore, by Definition 7.2, the vertex (v, u) is $X_i^{\pm}$-saturated. Thus C is $X_i^{\pm}$-saturated. Hence it is a cover of G_i.

By definition of precover, each nonempty connected component Φ of $\Gamma \times \Delta$ is a finite precover of G.

◇

Let C be the connected component of $\Gamma \times \Delta$ containing the vertex ϑ. Therefore $Lab(\Gamma \times \Delta, \vartheta) = Lab(C, \vartheta)$, because $Loop(\Gamma \times \Delta, \vartheta) = Loop(C, \vartheta)$. From now on we allow ourselves to vary between this two notations which define the same.

Lemma 7.4. *Let Γ and Δ be finite precovers of G such that $Lab(\Gamma, v) = H$ and $Lab(\Delta, u) = K$, where $v \in V(\Gamma)$ and $u \in V(\Delta)$.*

Let $\vartheta = (v, u) \in V(\Gamma \times \Delta)$. Let C be a connected component of $\Gamma \times \Delta$ such that $\vartheta \in V(C)$.

If $v \in VM_i(\Gamma)$ and $u \in VM_j(\Delta)$, where $1 \leq i \neq j \leq 2$, then

$$V(C) = \{\vartheta\}, \ E(C) = \emptyset \text{ and } \{1\} \leq H \cap K \leq A.$$

Otherwise $E(C) \neq \emptyset$ and $Lab(\Gamma \times \Delta, \vartheta) = H \cap K$.

Proof. If $v \in VM_i(\Gamma)$ and $u \in VM_j(\Delta)$, where $1 \leq i \neq j \leq 2$, then, by Definition 7.2, $V(C) = \{\vartheta\}$ and $E(C) = \emptyset$.

Let $H \cap K \neq \{1\}$. Then there exists $1 \neq w \in H \cap K$. Without loss of generality, we can assume that the word w is in normal form.

Assume first that the syllable length of the normal word w is greater than 1. Let $(x_1, x_2, \ldots, x_{k-1}, x_k)$ be a normal decomposition of w. By Lemma 6.1, this word labels a normal path closed at the basepoint in both graphs Γ and Δ.

Hence $x_1, x_k \in G_i \setminus A$, because $v \in VM_i(\Gamma)$. On the other hand, $u \in VM_j(\Delta)$ and therefore $x_1, x_k \in G_j \setminus A$ $(1 \leq i \neq j \leq 2)$. This is a contradiction. Therefore the syllable length of the normal word $w \in H \cap K$ is equal to 1. By Lemma 6.1, similar arguments show that in this case $w \in A$.

Assume now that the vertices v and u are not of different colors. Using the same ideas as in the proof of Lemma 7.3, we can assume, without loss of generality, that the vertices v and u are $X_i^{\pm}$-saturated $(i \in \{1, 2\})$. Therefore, by Definition 7.2, $E(C) \neq \emptyset$.

Let $w \in H \cap K$ be a normal word.

By Lemma 6.1, if either $l(w) > 1$, or $l(w) = 1$ and $w \notin A$ then the word w labels a path in Γ closed at v and also labels a path in Δ closed at u. Hence, by Definition 7.2, there exists a path p closed at ϑ in $C \subseteq \Gamma(H) \times \Gamma(K)$ such that $lab(p) \equiv w$. Thus $w \in Lab(\Gamma(H) \times \Delta, \vartheta) = Lab(C, \vartheta)$.

Assume now that the syllable length of w is equal to 1 and $w \in A$. Let $w' \in G_i \cap A$ such that $w' =_G w$. Since, by our assumption, the vertices v and u are $X_i^{\pm}$ saturated, Lemma 6.1 implies that the normal word w' labels a path closed at the basepoint in both graphs Γ and Δ. Therefore $w \in Lab(\Gamma(H) \times \Delta, \vartheta)$. Thus

$$H \cap K \subseteq Lab(\Gamma \times \Delta, \vartheta) = Lab(C, \vartheta).$$

Now let p be a path in $C \subseteq \Gamma \times \Delta$ closed at ϑ. Hence, by Definition 7.2, there exists a path p_1 in Γ closed at v and there exists a path p_2 in Δ closed at u, such that

$$lab(p) \equiv lab(p_1) \equiv lab(p_2).$$

Since $lab(p_1) \in Lab(\Gamma, v_0) = H$ and $lab(p_2) \in Lab(\Delta, u) = K$, we have $Lab(C, \vartheta) \subseteq Lab(\Gamma, v) \cap Lab(\Delta, u) = H \cap K$.

Hence $Lab(C, \vartheta) = H \cap K$.

◇

Remark 7.5. The above proof implies that if $H \cap K \leq G_i$ $(i \in \{1, 2\})$ then

$$H \cap K = Lab(C, v) \cap Lab(D, u),$$

where C and D are X_i-monochromatic component of Γ and Δ, respectively, such that $v \in V(C)$ and $u \in V(D)$. ◇

Recall that $(\Gamma(H), v_0)$ is the subgroup graph of H constructed by the generalized Stallings' algorithm.

Corollary 7.6. *Let H be a finitely generated subgroup of G.*

Let Δ be a finite precover of G such $Lab(\Delta, u) = K$.

Let $\vartheta = (v_0, u) \in V(\Gamma(H) \times \Delta)$. Let C be a connected component of $\Gamma(H) \times \Delta$ such that $\vartheta \in V(C)$.

Then $Lab(\Gamma(H) \times \Delta, \vartheta) = Lab(C, \vartheta) = H \cap K$.

Proof. If C is a nonempty (i.e., $E(C) \neq \emptyset$) connected component of $\Gamma(H)\times\Delta$ then, by Lemma 7.4, $Lab(\Gamma(H) \times \Delta, \vartheta) = Lab(C, \vartheta) = H \cap K$.

Otherwise, Lemma 7.4 implies that $\{1\} \leq H \cap K \leq A$. If $H \cap K = \{1\}$ then, since $Lab(C, \vartheta) = \{1\}$, the desired equality holds.

Assume now that $\{1\} \neq H\cap K \leq A$. Hence $H\cap A \neq \{1\}$. Therefore, since $\Gamma(H)$ is a reduced precover of G, Definition 5.4 (ii) implies $v \in VB(\Gamma(H))$. Therefore, by Lemma 7.4, $E(C) \neq \emptyset$. That is C is nonempty. This is a contradiction. Thus $Lab(\Gamma(H) \times \Delta, \vartheta) = Lab(C, \vartheta) = H \cap K$.

◇

Corollary 7.7 (The Intersection Problem)**.** *Let $H = \langle h_1, \cdots, h_n\rangle$ and $K = \langle k_1, \cdots, k_m\rangle$ be finitely generated subgroups of an amalgam of finite groups $G = G_1 *_A G_2$.*

Then there exists an algorithm which finds the generators of $H\cap K$, which is finitely generated.

Proof. We first use the generalized Stallings' folding algorithm to construct the subgroup graphs $(\Gamma(H), v_0)$ and $(\Gamma(K), u_0)$.

Since, by Theorem 4.1, these graphs are finite, the product graph $\Gamma(H)\times\Gamma(K)$ can be effectively constructed, and it is finite.

By Corollary 7.6, $H \cap K = Lab(C,(v_0, u_0))$, where C is a connected component of $\Gamma(H) \times \Gamma(K)$ such that $(v_0, u_0) \in V(C)$. Therefore it is sufficient to construct only the component C.

Let $\vartheta = (v_0, u_0)$. Recall that $G = G_1 *_A G_2 = gp\langle X \mid R\rangle$.

By Lemma 7.3, (C, ϑ) is a precover of G, which is finite by the construction. Therefore, in particular, it is a finite well-labelled graph. Hence, the subgroup $\widetilde{L}$ of $FG(X)$ determined by (C, ϑ) is finitely generated ([19, 29, 46]). Since $Lab(C, \vartheta) = \widetilde{L}/N = \widetilde{L}/\widetilde{L} \cap N$, where N is the normal closure of R in $FG(X)$, we conclude that $Lab(C, \vartheta) = H \cap K$ is finitely generated.

To find the generating set we proceed as follows. Let T be a fixed spanning tree of C. For all $v \in V(C)$, we consider t_v to be the unique freely reduced path in T from the basepoint ϑ to the vertex v.

For each $e \in E(C)$ we consider $t(e) = t_{\iota(e)}e\overline{t_{\tau(e)}}$. Thus if $e \in E(T)$ then $t(e)$ can be freely reduced to an empty path, that is $lab(t(e)) =_{FG(X)} 1$.

Let E^+ be the set of positively oriented edges of C. Let

$$X_L = \{lab(t(e)) \mid e \in E^+ \setminus E(T)\},$$

As is well known [19, 29, 46], $\widetilde{L} = FG(X_H)$. Therefore $L = \langle X_L \rangle$.

$\diamond$

Remark 7.8. In order to compute a finite group presentation of the subgroup $H \cap K$ one can apply to (C, ϑ) the restricted version of the Reidemeister-Schreier algorithm presented in [33]. This is possible, because C is a finite precover of G, by Lemma 7.3. $\diamond$

Corollary 7.9 (Howson Property)**.** *Let $G = G_1 *_A G_2$ be an amalgam of finite groups.*

The intersection of two finitely generated subgroups of G is finitely generated in G. That is G possesses the *Howson property.*

Malnormality.

Lemma 7.10. *Let H and K be finitely generated subgroups of the group G. Let g be an element of G.*

Then $H \cap gKg^{-1}$ conjugates to a subgroup of A or it conjugates to the subgroup $Lab(C, \vartheta)$, where C is a nonempty connected component of $\Gamma(H) \times \Gamma(K)$ such that $\vartheta = (v, u) \in V(C)$, and $v \in V(\Gamma(H))$, $u \in V(\Gamma(K))$.

Proof. Without loss of generality, assume that g is a normal word. Then either there exists a path p in $\Gamma(K)$ such that $\iota(p) = u_0$ and $lab(p) \equiv g^{-1}$ or such a path doesn't exist in $\Gamma(K)$.

In the first case, let $u = \tau(p)$ then $Lab(\Gamma(K), u) = gKg^{-1}$. By Corollary 7.6, $Lab(C, \vartheta) = H \cap gKg^{-1}$, where $\vartheta = (v_0, u)$.

Assume now that p' is the longest path in $\Gamma(K)$ such that $\iota(p') = u_0$ and $lab(p') \equiv g_2^{-1}$, where $g \equiv g_1 g_2$. Let $u = \tau(p')$. Then either there exists a path q in $\Gamma(H)$ such that $\iota(q) = v_0$ and $lab(q) \equiv g_1$ or such a path doesn't exist in $\Gamma(H)$.



Figure 9.

First we assume that q exists in $\Gamma(H)$, see Figure 9. Let $v = \tau(q)$. Thus $Lab(\Gamma(K), u) = g_2 K g_2^{-1}$ and $Lab(\Gamma(H), v) = g_1^{-1} H g_1$. By Lemma 7.4, if u

and v are monochromatic vertices of different colors then

$$\{1\} \leq g_1^{-1}Hg_1 \cap g_2Kg_2^{-1} \leq A.$$

Otherwise, $Lab(C,(u,v)) = g_1^{-1}Hg_1 \cap g_2Kg_2^{-1}$, where C is a nonempty connected component of the product graph $\Gamma(H) \times \Gamma(K)$ containing the vertex $\vartheta = (v,u)$, $v \in V(\Gamma(H))$, $u \in V(\Gamma(K))$.

Since

$$H \cap gKg^{-1} = g_1(g_1^{-1}Hg_1 \cap g_2Kg_2^{-1})g_1^{-1},$$

we have $\{1\} \leq H \cap gKg^{-1} \leq g_1Ag_1^{-1}$ or $H \cap gKg^{-1} = g_1Lab(C,\vartheta)g_1^{-1}$, respectively.

Assume now that there is no path in $\Gamma(H)$ starting at v_0 and labelled with g_1. Below we prove that in this case $H \cap gKg^{-1} = \{1\}$.

Suppose that $H \cap gKg^{-1} \neq \{1\}$. Let (Γ', u') be the graph obtained from $\Gamma(K)$ by attaching a path t at the vertex u, such that $\iota(t) = u$ and $lab(t) \equiv g_1^{-1}$. Let $\tau(t) = u'$, see Figure 10. The graph (Γ', u') is finite, because $\Gamma(K)$ is finite and the "stem" t is also finite. It is well-labelled, because $\Gamma(K)$ is well-labelled and g_2^{-1} is the maximal prefix of the word g^{-1} that is readable in $\Gamma(K)$ starting at u_0.



FIGURE 10. The graph Γ'

Thus $Lab(\Gamma', u') = gKg^{-1}$, and for each nontrivial element in gKg^{-1} and ,in particular, for each $1 \neq z \in H \cap gKg^{-1}$ there exists a nonempty path γ in (Γ', u') such that $lab(\gamma) =_G z$ and $\iota(\gamma) = \tau(\gamma) = u'$. The above construction of (Γ', u') implies that $lab(\gamma) \equiv g_1wg_1^{-1}$, where $w \in Lab(\Gamma(K), u)$. Since $lab(\gamma) =_G z \neq 1$, the word w is nonempty. Thus, by Lemma 6.1, we can assume that the word w is in normal form, because $(\Gamma(K), u)$ is a finite precover of G and $w \in Lab(\Gamma(K), u)$.

Since each X_i-monochromatic component of $\Gamma(K)$ is a cover of G_i, $i \in \{1,2\}$ (thus, in particular, it is $X_i^{\pm}$-saturated) and because g_2^{-1} is the maximal prefix of the word g^{-1} such that there is a path p' in $\Gamma(K)$ with $\iota(p') = u_0$ and $lab(p') \equiv g_2^{-1}$, there exists a normal decomposition of the word g

$$(y_1, \ldots, y_k, x_1, \ldots, x_m)$$

such that $g_1 \equiv y_1 \cdots y_k$ and $g_2 \equiv x_1 \cdots x_m$, where $y_k \in G_i$ and $x_1 \in G_j$, $1 \leq i \neq j \leq 2$.

Note that u is a X_j-monochromatic vertex of $\Gamma(K)$. Otherwise there exists a G_i-monochromatic component D in $\Gamma(K)$, such that $u \in V(D)$. Since it is X_i-saturated, y_k is readable from u in D and therefore in $\Gamma(K)$. This contradicts the maximality of the word g_2. Thus the word $g_1 w g_1^{-1}$ is in normal form.

On the other hand, since $z =_G g_1 w g_1^{-1} \in H$, Theorem 4.1 (4) implies that there exists a normal path γ' in $\Gamma(H)$ closed at v_0 with $lab(\gamma') \equiv g_1 w g_1^{-1}$. Therefore there exists a path in $\Gamma(H)$ starting at v_0 and labelled with g_1. This contradicts with our assumption that such a path doesn't exist in $\Gamma(H)$. Hence $H \cap gKg^{-1} = \{1\}$.

◇

Theorem 7.11. *Let H be a finitely generated subgroup of G. Then H is malnormal in G if and only if the following holds*

1. $H \cap gHg^{-1} \cap fAf^{-1} = \{1\}$ *for all* $g \in G \setminus H$, $f \in G$;
2. *each connected nonempty component C of $\Gamma(H) \times \Gamma(H)$ which doesn't contain the vertex (v_0, v_0) satisfies $Lab(C, \vartheta) = \{1\}$ for all $\vartheta \in V(C)$.*

Proof. Suppose that H is malnormal in G. Then $H \cap gHg^{-1} = \{1\}$ for all $g \in G \setminus H$. Hence $H \cap gHg^{-1} \cap fAf^{-1} = \{1\}$ for all $g, f \in G$.

Let C be a nonempty connected component of $\Gamma(H) \times \Gamma(H)$ such that $(v_0, v_0) \notin V(C)$. Let $\vartheta = (v_1, v_2) \in V(C)$. Hence $v_1 \neq v_2 \in V(\Gamma(H))$. Indeed, if $v_1 = v_2$ then $(v_1, v_2) \in V(C_0)$, where C_0 is a connected component of the product graph $\Gamma(H) \times \Gamma(H)$, containing the vertex (v_0, v_0).

Lemma 7.4 implies that

$$Lab(C, \vartheta) = Lab(\Gamma(H), v_1) \cap Lab(\Gamma(H), v_2) = g_1^{-1} H g_1 \cap g_2^{-1} H g_2,$$

where g_1 and g_2 label paths in $\Gamma(H)$ from v_0 to v_1 and to v_2, respectively.

Since $\Gamma(H)$ is a G-based graph and $v_1 \neq v_2$, we have $g_1 g_2^{-1} \notin H$. Indeed, otherwise

$$v_0 = v_0 \cdot (g_1 g_2^{-1}) = (v_0 \cdot g_1) \cdot g_2^{-1} = v_1 \cdot g_2^{-1}.$$

Thus

$$v_1 = v_1 \cdot (g_2^{-1} g_2) = (v_1 \cdot g_2^{-1}) \cdot g_2 = v_0 \cdot g_2 = v_2.$$

However $g_1 Lab(C, \vartheta) g_1^{-1} = H \cap g_1 g_2^{-1} H g_2 g_1^{-1} = \{1\}$, because H is malnormal in G. Therefore $Lab(C, \vartheta) = \{1\}$.

Assume now that the conditions (1)-(2) are satisfied. By Lemma 7.10, the subgroup $H \cap gHg^{-1}$ conjugates to a subgroup of A or it conjugates to the subgroup $Lab(C, \vartheta)$, where C is a nonempty (i.e. $E(C) \neq \emptyset$) connected component of the product graph $\Gamma(H) \times \Gamma(H)$.

In the first case, $1 \leq f^{-1}(H \cap gHg^{-1})f \leq A$ for some $f \in G$. Therefore $H \cap gHg^{-1} \cap fAf^{-1} \neq \{1\}$. This contradicts condition (1).

Condition (2) implies $Lab(C, \vartheta) = \{1\}$, hence $H \cap gHg^{-1} = \{1\}$. Therefore H is malnormal in G.

◇

Corollary 7.12 (The Malnormality Problem). *Let $h_1, \dots h_k \in G$. Then there exists an algorithm which decides whether or not the subgroup $H = \langle h_1, \dots h_k \rangle$ is malnormal in G.*

If H is not malnormal, the algorithm produces a nontrivial element $g \in G \setminus H$ such that $H \cap gHg^{-1} \neq \{1\}$.

Proof. First we construct the subgroup graph $\Gamma(H)$ using the generalized Stallings' algorithm. Since, by Theorem 4.1, it is finite, the product graph $\Gamma(H) \times \Gamma(H)$ can be constructed effectively. Now we check whether each connected nonempty component C of $\Gamma(H) \times \Gamma(H)$ which doesn't contain the vertex (v_0, v_0) satisfies $Lab(C, \vartheta) = \{1\}$ for some $\vartheta \in V(C)$. If there exists a component with $Lab(C, \vartheta) \neq \{1\}$ ($\vartheta \in V(C)$), then, by Theorem 7.11, H is not malnormal in G.

Moreover, by the proof of Theorem 7.11, a nontrivial element $g \in G \setminus H$ such that $H \cap gHg^{-1} \neq \{1\}$ is $g =_G g_1 g_2^{-1}$, where

$$v_1 = v_0 \cdot g_1, \quad v_2 = v_0 \cdot g_2 \ \text{ and } \ \vartheta = (v_1, v_2).$$

Note that it is sufficient to check whether $Lab(C, \vartheta) = \{1\}$ only for some $\vartheta \in V(C)$. Indeed, if $v' \in V(C)$ such that $v' \neq \vartheta$ then $Lab(C, v') = xLab(\Gamma, \vartheta)x^{-1}$, where $x \in G$ and $v' \cdot x = \vartheta$. Thus $Lab(\Gamma, v') = \{1\}$.

Since, by Lemma 7.3, C is a precover, the above verification can be done as follows.

By Lemma 5.7(i), a reduced precover (Δ, u) has $Lab(\Delta, u) = \{1\}$ if and only if $V(\Delta) = u$ and $E(\Delta) = \emptyset$. Thus $Lab(C, \vartheta) = \{1\}$ if and only if the iterative removal of the unique sequence of redundant components from (C, ϑ) yield the empty graph (Δ, u) with the above properties.

Assume now that all connected nonempty components of $\Gamma(H) \times \Gamma(H)$ satisfy condition (2) from Theorem 7.11. Then H is malnormal in G if and only if condition (1) is satisfied. In order to verify this we proceed as follows.

Let D be an arbitrary single vertex of the product graph $\Gamma(H) \times \Gamma(H)$, i.e. D is an empty component of $\Gamma(H) \times \Gamma(H)$ such that $V(D) = \{(v_1, v_2)\}$ and $E(D) = \emptyset$. Then $v_1 \neq v_2 \in V(\Gamma(H))$ such that $v_0 \cdot g_i = v_i$, $i \in \{1, 2\}$. Since $E(D) = \emptyset$, by Lemma 7.4, v_1 and v_2 are monochromatic vertices of $\Gamma(H)$ of different colors. Without loss of generality, assume that $v_i \in VM_i(\Gamma(H))$, $i \in \{1, 2\}$. By Lemma 7.4, $\{1\} \leq g_1^{-1} H g_1 \cap g_2^{-1} H g_2 \leq A$.

Let C_i be a X_i-monochromatic component of $\Gamma(H)$ such that $v_i \in V(C_i)$. By Remark 7.5,

$$g_1^{-1}Hg_1 \cap g_2^{-1}Hg_2 = Lab(C_1, v_1) \cap Lab(C_2, v_2) \neq \{1\}.$$

Thus we have to check if $Lab(C_1, v_1) \cap Lab(C_2, v_2)$ is a nontrivial subgroup of A. If so then $g_1^{-1}Hg_1 \cap g_2^{-1}Hg_2$ is a nontrivial subgroup of A, otherwise $g_1^{-1}Hg_1 \cap g_2^{-1}Hg_2 = \{1\}$.

Let $S = A \cap Lab(C_1, v_1)$. We consider $(Cayley(G_2, S), S \cdot 1)$. Thus $Lab(Cayley(G_2, S), S \cdot 1) = S$. Let E be a nonempty connected component of the product graph $Cayley(G_2, S) \times C_2$ containing the vertex $(S \cdot 1, v_2)$. Then, by Lemma 7.4,

$$Lab(E, (S \cdot 1, v_2)) = Lab(Cayley(G_2, S), S \cdot 1) \cap Lab(C_2, v_2) =$$

$$= S \cap Lab(C_2, v_2) = A \cap Lab(C_1, v_1) \cap Lab(C_2, v_2) = g_1^{-1}Hg_1 \cap g_2^{-1}Hg_2.$$

Thus $g_1 Lab(E, (S \cdot 1, v_2))g_1^{-1} = H \cap gHg^{-1}$, where $g =_G g_1g_2^{-1}$. Hence $Lab(E, (S \cdot 1, v_2)) \neq \{1\}$ implies H is not malnormal in G.

Otherwise if $Lab(E, (S \cdot 1, v_2)) = \{1\}$ for each component D of the product graph $\Gamma(H) \times \Gamma(H)$, where E is constructed as described above, then H is a malnormal subgroup of G.

◇

Example 7.13. Let $G = gp\langle x, y | x^4, y^6, x^2 = y^3 \rangle = G_1 *_A G_2$, where $G_1 = gp\langle x | x^4 \rangle$, $G_2 = gp\langle y | y^6 \rangle$ and $A = \langle x^2 \rangle = \langle y^3 \rangle$.

Let H be a finitely generated subgroup of G given by its subgroup graph $\Gamma(H)$ which is presented on Figure 11.

We compute $\Gamma(H) \times \Gamma(H)$ (see Figure 11). Using the method described along with the proof of Corollary 7.12, we conclude that $Lab(C_1, (v_0, v_1)) = \{1\}$, $Lab(C_3, (v_0, v_3)) = \{1\}$, but $Lab(C_2, (v_0, v_2)) \neq \{1\}$. Therefore, by Theorem 7.11, H is not malnormal in G.

◇

Complexity. By Theorem 4.1 (5), the complexity of the construction of $\Gamma(H)$ for a subgroup H of G given by a finite set of generators is $O(m^2)$, where m is the sum of lengths of the input subgroup generators.

The construction of $\Gamma(H) \times \Gamma(H)$ takes

$$O(|V(\Gamma(H))|^2 + |V(\Gamma(H))| \cdot |E(\Gamma(H))|).$$

Let C be a connected component of $\Gamma(H) \times \Gamma(H)$. To verify whether $Lab(C, v) = 1$, $v \in V(C)$, takes time proportional to $|E(C)|^2$, by the complexity analysis of the generalized Stallings algorithm (see Lemma 8.7 in



FIGURE 11.

[32]). Since

$$\sum_{C\subseteq\Gamma(H)\times\Gamma(H)} |E(C)| = |E(\Gamma(H)\times\Gamma(H))| \leq |E(\Gamma(H))|^2,$$

the above verification for all connected components of $\Gamma(H)\times\Gamma(H)$ takes $O(|E(\Gamma(H))|^4)$.

Since all the information about the free factors of the amalgams, as well as the relative Cayley graphs of the free factors are not a part of the input, the verifications concerning the empty components of the product graph $\Gamma(H)\times\Gamma(H)$ takes time $O(|V(\Gamma)|^2)$.

Since, by Theorem 4.1 (5), $|E(\Gamma(H))|$ and $|V(\Gamma(H))|$ are proportional to m, algorithm given by the proof of Corollary 7.12 takes $O(m^4)$. Thus the algorithm is polynomial in the size of the input.

Almost Malnormality.

Theorem 7.14. *Let H be a finitely generated subgroup of G. Then H is almost malnormal in G if and only if $Lab(C,\vartheta)$ conjugates to a subgroup of G_1 or G_2 ($\vartheta\in V(C)$), for each nonempty connected component C of $\Gamma(H)\times\Gamma(H)$, which doesn't contain the vertex (v_0,v_0).*

Proof. Suppose that H is almost malnormal in G. Then $H\cap gHg^{-1}$ is finite for all $g\in G\setminus H$.

Let C be a nonempty connected component of $\Gamma(H)\times\Gamma(H)$ such that $(v_0,v_0)\notin V(C)$. Let $\vartheta=(v_1,v_2)\in V(C)$. By the proof of Theorem 7.11, $v_1\neq v_2\in V(\Gamma(H))$ and $g_1g_2^{-1}\notin H$, where $v_i=v_0\cdot g_i$ ($i\in\{1,2\}$). Moreover,

$$Lab(C,\vartheta) = Lab(\Gamma(H),v_1)\cap Lab(\Gamma(H),v_2) = g_1^{-1}Hg_1\cap g_2^{-1}Hg_2.$$

However $g_1 Lab(C, \vartheta) g_1^{-1} = H \cap g_1 g_2^{-1} H g_2 g_1^{-1}$ is finite, because H is almost malnormal in G. Therefore, by the Torsion Theorem (IV.2.7 in [27]), $g_1 Lab(C, \vartheta) g_1^{-1}$ conjugates to a subgroup of G_1 or G_2.

Assume now that the condition is satisfied. By Lemma 7.10, for all $g \in G \backslash H$ the subgroup $H \cap gHg^{-1}$ conjugates to a subgroup of A or it conjugates to the subgroup $Lab(C, \vartheta)$, where C is a nonempty connected component of the product graph $\Gamma(H) \times \Gamma(H)$. Therefore if $Lab(C, \vartheta)$ conjugates to a subgroup of G_1 or G_2, then, since G_i $(i \in \{1, 2\})$ is finite, H is almost malnormal.

◇

The *almost malnormality problem* asks to decide whether or not a subgroup H of the group G is almost malnormal in G.

Corollary 7.15 (The Almost Malnormality Problem). *Let $h_1, \ldots h_k \in G$. Then there exists an algorithm which decides whether or not the subgroup $H = \langle h_1, \ldots h_k \rangle$ is almost malnormal in G.*

If H is not almost malnormal, the algorithm produces a nontrivial element $g \in G \setminus H$ such that $H \cap gHg^{-1}$ is not finite.

Proof. The proof is similar to that of Corollary 7.12. First we construct the subgroup graph $\Gamma(H)$ using the generalized Stallings' algorithm. Since, by Theorem 4.1, it is finite, the product graph $\Gamma(H) \times \Gamma(H)$ can be constructed effectively. Now for each nonempty connected component C of $\Gamma(H) \times \Gamma(H)$ which doesn't contain the vertex (v_0, v_0), we check whether $Lab(C, \vartheta)$ $(\vartheta \in V(C))$ conjugates to a subgroup of G_1 or G_2. By Theorem 7.14, H is almost malnormal in G if and only if each such component C possesses this property.

We proceed as follows. If C consists of a unique X_i-monochromatic component $(i \in \{1, 2\})$ then $\{1\} \leq Lab(C, v) \leq G_i$. Otherwise, let $\vartheta \in VB(C)$ be a basepoint of C.

By Lemma 7.3, C is a finite precover of G. If (C, ϑ) is not a reduced precover then we remove from C all the redundant components w.r.t. the basepoint ϑ. Let (C', ϑ') be the resulting graph, where ϑ' is the image of ϑ in C'. Thus (C', ϑ') is a reduced precover such that $Lab(C, \vartheta) = Lab(C', \vartheta')$. Let $L = Lab(C', \vartheta')$.

By Lemma 5.7, $L \leq G_i$ such that $L \cap A = \{1\}$ if and only if C' consists of a unique X_i-monochromatic component, and $L \leq A$ if and only if $(C', \vartheta') = Cayley(G_1, L) *_{\{La \mid a \in A\}} Cayley(G_2, L)$. Thus $Lab(C, \vartheta) = L$ conjugates to a subgroup of G_1 or G_2 if and only if C' satisfies one of the above properties.

Note that if there exists a connected component C such that none of the above properties is satisfied then, by the proof of Theorem 7.14, a nontrivial element $g \in G \setminus H$ such that $H \cap gHg^{-1}$ is not finite is $g =_G g_1 g_2^{-1}$, where

$$v_1 = v_0 \cdot g_1, \quad v_2 = v_0 \cdot g_2 \text{ and } \vartheta = (v_1, v_2).$$

$\diamond$

Complexity. Similarly to the complexity analysis of the solution of the malnormality problem, presented along with the proof of Corollary 7.12, the above solution of the almost malnormality problem takes $O(|E(\Gamma(H))|^4)$, that is $O(m^4)$, where $m = \sum_{i=1}^{k} |h_i|$.

8. The Power Problem

The *power problem* asks for an algorithm that decides whether or not some *nontrivial power* of a word g in the generators of a group G belongs to the subgroup H of G.

By a *nontrivial power* of g we mean an element $g^n \in G$ such that $n \geq 1$ and $g^n \neq_G 1$ (otherwise $g^n \in H$ for each torsion element $g \in G$ and all $o(g) \mid n$).

This problem is an extension of the membership problem for H in G. The membership problem for finitely generated subgroups in amalgams of finite groups was (successfully) solved in [32] using subgroup graphs constructed by the generalized Stallings' algorithm. Below we employ same technics to solve the power problem in this class of groups (Corollaries 8.1 and 8.4). Theorem 8.3 provides the solution. The complexity analysis of the described algorithm is given at the end of the section.

We split the power problem into two instances. The first one, (PI), asks for an answer "Yes" or "No" on the question whether some nonzero power of a word g in the generators of G belongs to the subgroup H.

The second one, (PII), asks to find the minimal power $n > 0$ such that $g^n \in H$. Evidently, (PII) implies (PI).

Corollary 8.1 (The Power Problem)**.** *Let $G = G_1 *_A G_2$ be an amalgam of finite groups. Then there exists an algorithm which solves (PI).*

That is, given finitely many subgroup generators $h_1, \dots h_k \in G$ and normal word $g \in G$, the algorithm decides whether or not some nonzero power of g is in the subgroup $H = \langle h_1, \dots h_k \rangle$.

Proof. Let $K = \langle g \rangle$. Construct the subgroup graphs $(\Gamma(H), v_0)$ and $(\Gamma(K), u_0)$ using the generalized Stallings' algorithm.

By Corollary 7.6, $Lab(C, \vartheta) = H \cap K = \langle g^n \rangle$, where C is the connected component of $\Gamma(H) \times \Gamma(K)$ such that $\vartheta = (v_0, u_0) \in V(C)$. Therefore $Lab(C, \vartheta) = \{1\}$ implies no nonzero power of g is in H.

Thus we construct the connected component C of the product graph $\Gamma(H) \times \Gamma(K)$. The verification whether or not $Lab(C, \vartheta) = \{1\}$ can be done as is explained in the proof of Corollary 7.12.

◇

Complexity. By the complexity analysis of the "malnormality" algorithm given along with the proof of Corollary 7.12, the complexity of the above algorithm given by Corollary 8.1 is $O\big(|E(\Gamma(H))|^2 \cdot |E(\Gamma(K))|^2\big)$. That is $O\big(m^2 \cdot |g|^2\big)$, where m is the sum of lengths of $h_1, \dots h_k$.

Following [27], we say that a word $g \equiv g_1 g_2 \cdots g_k \in G = G_1 *_A G_2$ given by the normal decomposition $(g_1, g_2, \dots, g_k)$ is *cyclically reduced* if $k \leq 1$ or if g_1 and g_k are in different factors of G. Hence if g is cyclically reduced then all cyclic permutations of $(g_1, g_2, \dots, g_k)$ define normal words. Obviously, if $g \in G$ is cyclically reduced then g is *freely cyclically reduced*, that is $g \not\equiv x g' x^{-1}$ $(x \in X^{\pm})$.

Lemma 8.2. *Let $g \in G$ be a normal word given by the normal decomposition $(g_1, g_2, \dots, g_k)$. Then there exists a normal word $x \in G$ and a cyclically reduced word $g' \in G$ such that $g =_G x g' x^{-1}$ and the word $x g' x^{-1}$ is in normal form.*

Proof. If $k = 1$ then the statement is trivial: $x \equiv 1$ and $g' \equiv g$. If k is an even number then the syllables g_1 and g_k are, evidently, in different free factors. Therefore g is cyclically reduced. Thus the statement is trivial.

Assume now that k is odd. The proof is by induction on the syllable length of g, that is on k.

If $g_k g_1 \in G_i \setminus A$ then we put $g'_k =_{G_i} g_k g_1$ $(i \in \{1, 2\})$. Thus $g =_G g_1 (g_2 \cdots g'_k) g_1^{-1}$. Since $g_2 \in G_j \setminus A$ $(1 \leq i \neq j \leq 2)$, the word $g' \equiv g_2 \cdots g'_k$ is normal and cyclically reduced. Moreover, the words $x \equiv g_1$ and $x g' x \equiv g_1 g_2 \cdots g_{k-1} (g'_k g_1^{-1})$ are normal.

If $g_k g_1 \in G_i \cap A$ $(i \in \{1, 2\})$ we take $b \in G_j \cap A$ $(1 \leq i \neq j \leq 2)$ such that $b =_G g_k g_1$. Since $g_{k-1} \in G_j \setminus A$, we have $g_{k-1} b \in G_j \setminus A$. Let $g'_{k-1} =_{G_j} g_{k-1} b$. Then

$$g =_G g_1 (g_2 \cdots g_{k-2} g'_{k-1}) g_1^{-1}.$$

We put $x \equiv g_1$, $g' \equiv g_2 \cdots g_{k-2} g'_{k-1}$. Thus x, g' and $x g' x^{-1}$ are normal words. Moreover, by the inductive assumption, there exists a normal word $x' \in G$ and a cyclically reduced word $g'' \in G$ such that $g' =_G x' g'' (x')^{-1}$, where $x' \equiv g_2 \cdots g_l$ $(l \leq (n-1)/2)$ and the word $x' g'' (x')^{-1}$ is in normal form.

Thus the words $x x'$ and $(x x') g'' (x x')^{-1}$ are normal. Since $g =_G (x x') g'' (x x')^{-1}$, we are done.

◇

Theorem 8.3. *Let H be a finitely generated subgroup of an amalgam of finite groups $G = G_1 *_A G_2$. Let $g \in G$ be a non torsion element such that $g^n \in H$ for some $n \geq 1$.*

Then there exists $1 \leq z \leq |V(\Gamma(H))|$ such that $g^z \in H$.

Proof. Assume that $g \notin H$ otherwise the statement is trivial. Let $n \geq 1$ be the smallest positive integer such that $g^n \in H$. Since $g \notin H$, we have $n > 1$. Suppose that $n > |V(\Gamma(H))| > 1$ otherwise the statement is trivial.

Without loss of generality we can assume that g is a normal word given by the normal decomposition $g \equiv g_1 \cdots g_k$, where $k > 1$ since g is non torsion.

By Lemma 8.2, there exits a normal word $x \in G$ and a cyclically reduced word $g' \in G$, such that $g =_G xg'x^{-1}$ and the word $xg'x^{-1}$ is normal.

Note that the syllable length of g' is greater than 1. Otherwise g' is an element of either G_1 or of G_2. Thus g is a conjugate of an element of either G_1 or of G_2. Therefore, by the Torsion Theorem (IV.2.7 in [27]), g is a torsion element of G, which contradicts our assumption.

Therefore $g^n =_G x(g')^n x^{-1}$ and the word $x(g')^n x^{-1}$ is normal. Hence, by Theorem 4.1 (4), there exists a normal path p in $\Gamma(H)$ with $\iota(p) = \tau(p) = v_0$ and $lab(p) \equiv x(g')^n x^{-1}$. Since the graph $\Gamma(H)$ is well-labelled, there is a decomposition $p = tq\overline{t}$, where

$$\iota(t) = v_0, \ \tau(t) = v, \ lab(t) \equiv x, \ \text{and} \ \iota(q) = \tau(q) = v, lab(q) \equiv (g')^n.$$

Since the word g' is freely cyclically reduced, we have $|(g')^n| = |g'| \cdot n$. Hence we can set $v_m = u \cdot (g')^m$, $1 \leq m \leq n$. Since $n > |V(\Gamma(H))|$, there exist $1 \leq i < j \leq n$ such that $v_i = v_j$. Thus $v_i \cdot (g')^{j-i} = v_j = v_i$. Therefore $v = v \cdot (g')^n = v \cdot (g'v)^{n-(j-i)}$. Hence

$$v_0 \cdot \left(x(g')^{n-(j-i)}x^{-1}\right) = v \cdot \left((g')^{n-(j-i)}x^{-1}\right) = v \cdot x^{-1} = v_0.$$

Thus $x(g')^{n-(j-i)}x^{-1} \in H$. Hence $g^{n-(j-i)} \in H$. Since $1 \leq i < j$, we have $1 \leq n - (j - i) < n$. This contradicts with the choice of n.

◇

Corollary 8.4 (The Power Problem)**.** *Let $G = G_1 *_A G_2$ be an amalgam of finite groups. Then there exists an algorithm which solves (PII).*

That is, given finitely many subgroup generators $h_1, \ldots h_k \in G$ and normal word $g \in G$, the algorithm finds the minimal nonzero power n such that $g^n \in H = \langle h_1, \ldots h_k \rangle$.

Proof. We begin by rewriting the word g as a normal word $xg'x^{-1}$, where $x \in G$ is a normal word and $g' \in G$ is a cyclically reduced word. This is possible by Lemma 8.2 and can be done according to the process described in the proof of this lemma. Thus $g =_G xg'x^{-1}$.

If $l(g') = 1$ then $g' \in G_i$ ($i \in \{1, 2\}$). Thus g is a torsion element of G. Let $o(g')$ be the order of g'. Since $1 < o(g) = o(g') \leq |G_i|$, we have to verify whether $g^m \in H$, for all $1 \leq m \leq |G_i| - 1$, and to stop when the first such power is found or when $g^m =_G 1$, that is no nontrivial power of g is in H.

By Theorem 4.1 (4), such a verification can be done using the subgroup graph $(\Gamma(H), v_0)$ constructed by the generalized Stallings' algorithm. That is $g^m \in H$ if and only if its normal form labels a normal path in $\Gamma(H)$ closed at the basepoint v_0. If $(g')^m \notin A$ $(1 \leq m \leq |G_i| - 1)$, then $x(g')^m x$ is a normal word. Otherwise we just rewrite it as a normal word.

If $l(g') > 1$, then, by the proof of Theorem 8.3, $g^m \in H$ if and only if there exists a path p in $\Gamma(H)$ closed at v_0 with $lab(p) \equiv x(g')^m x^{-1}$ such that $1 \leq m \leq |V(\Gamma(H))|$.

Hence we try to read $x(g')^m x^{-1}$ on $\Gamma(H)$ starting at v_0, for all $1 \leq m \leq |V(\Gamma(H))|$. That is we begin with $m = 1$ and stop when we succeed to read $x(g')^m x^{-1}$ at the first time. If no such m is found then no nonzero power of g is in H.

◇

Complexity. By Theorem 4.1 (5), the construction of $\Gamma(H)$ takes $O(m^2)$, where m is the sum of the lengths of $h_1, \dots h_k$. To find the desired normal form of g, which is $xg'x^{-1}$, takes $O(|g|)$. A verification of whether or not $x(g')^i x^{-1}$ can be read on $\Gamma(H)$ starting at v_0 $(1 \leq i \leq |V(\Gamma(H)|)$ takes $O(|g| \cdot |V(\Gamma(H)|)$, when g is non torsion. Otherwise it takes $O(|g| \cdot |G_i|)$ ($i \in \{1, 2\}$). Since the information about the factors, G_1 and G_2, is given and it is not a part of the input, it takes $O(|g|)$.

Since, by Theorem 4.1 (5), $|V(\Gamma(H))|$ is proportional to m, we conclude that the complexity of the algorithm given along with the proof of Corollary 8.4 is $O(m^2 + m \cdot |g|)$. Thus the algorithm is quadratic in the size of the input. Moreover, it is faster than the algorithm presented in Corollary 8.1 which solves (PI).

Appendix A.

Below we follow the notation of Grunschlag [16], distinguishing between the "*input*" and the "*given data*", the information that can be used by the algorithm *"for free"*, that is it does not affect the complexity issues.

Algorithm

Given: Finite groups G_1, G_2, A and the amalgam $G = G_1 *_A G_2$ given via (1.a), (1.b) and (1.c), respectively.

We assume that the Cayley graphs and all the relative Cayley graphs of the free factors are given.

Input: A finite set $\{g_1, \cdots, g_n\} \subseteq G$.

Output: A finite graph $\Gamma(H)$ with a basepoint v_0 which is a reduced precover of G and the following holds

- $Lab(\Gamma(H), v_0) =_G H$;
- $H = \langle g_1, \cdots, g_n \rangle$;
- a normal word w is in H if and only if there is a loop (at v_0) in $\Gamma(H)$ labelled by the word w.

Notation: Γ_i is the graph obtained after the execution of the i-th step.

Step1: Construct a based set of n loops around a common distinguished vertex v_0, each labelled by a generator of H;

Step2: Iteratively fold edges and cut hairs [8];

Step3:

```
For each $X_i$-monochromatic component $C$ of $\Gamma_2$ $(i = 1, 2)$ Do
Begin
pick an edge $e \in E(C)$;
glue a copy of $Cayley(G_i)$ on $e$ via identifying $1_{G_i}$ with $\iota(e)$
and identifying the two copies of $e$ in $Cayley(G_i)$ and in $\Gamma_2$;
If necessary Then iteratively fold edges;
End;
```

Step4:

```
For each $v \in VB(\Gamma_3)$ Do
If there are paths $p_1$ and $p_2$, with $\iota(p_1) = \iota(p_2) = v$ and
$\tau(p_1) \neq \tau(p_2)$ such that
    $lab(p_i) \in G_i \cap A$ $(i = 1, 2)$ and $lab(p_1) =_G lab(p_2)$
 Then identify $\tau(p_1)$ with $\tau(p_2)$;
If necessary Then iteratively fold edges;
```

Step5: Reduce Γ_4 by an iterative removal of all (*redundant*) X_i-monochromatic components C such that

- (C, ϑ) is isomorphic to $Cayley(G_i, K, K \cdot 1)$, where $K \leq A$ and $\vartheta \in VB(C)$;
- $|VB(C)| = [A : K]$;
- one of the following holds
  - $K = \{1\}$ and $v_0 \notin VM_i(C)$;
  - K is a nontrivial subgroup of A and $v_0 \notin V(C)$.

Let Γ be the resulting graph;

---

[8] A *hair* is an edge one of whose endpoint has degree 1

```
    If     $VB(\Gamma) = \emptyset$ and $(\Gamma, v_0)$ is isomorphic to $Cayley(G_i, 1_{G_i})$
    Then   we set $V(\Gamma_5) = \{v_0\}$ and $E(\Gamma_5) = \emptyset$;
    Else   we set $\Gamma_5 = \Gamma$.
Step6:
  If
    • $v_0 \in VM_i(\Gamma_5)$ $(i \in \{1, 2\})$;
    • $(C, v_0)$ is isomorphic to $Cayley(G_i, K, K \cdot 1)$, where $L = K \cap A$
      is a nontrivial subgroup of $A$ and $C$ is a $X_i$-monochromatic
      component of $\Gamma_5$ such that $v_0 \in V(C)$;
  Then   glue to $\Gamma_5$ a $X_j$-monochromatic component $(1 \leq i \neq j \leq 2)$
  $D = Cayley(G_j, L, L \cdot 1)$ via identifying $L \cdot 1$ with $v_0$ and
  identifying the vertices $L \cdot a$ of $Cayley(G_j, L, L \cdot 1)$ with the vertices
  $v_0 \cdot a$ of $C$, for all $a \in A \setminus L$.
    Denote $\Gamma(H) = \Gamma_6$.
```

Remark A.1. Note that the first two steps of the above algorithm correspond precisely to the Stallings' folding algorithm for finitely generated subgroups of free groups [46, 29, 19]. ◇



FIGURE 12. The construction of $\Gamma(H_1)$.

Example A.2. Let $G = gp\langle x, y | x^4, y^6, x^2 = y^3 \rangle$.

Let H_1 and H_2 be finitely generated subgroups of G such that

$$H_1 = \langle xy \rangle \text{ and } H_2 = \langle xy^2, yxyx \rangle.$$

The construction of $\Gamma(H_1)$ and $\Gamma(H_2)$ by the algorithm presented above is illustrated on Figures 12 and 13. ◇

References

[1] M. Bestvina, M. Feighn, A combinatorial theorem for negatively curved groups, *J. Differential Geom.* **35** (1992), no.1, 85-101.

[2] M. Bestvina, M. Feighn, Addendum and correction to: "A combination theorem for negatively curved groups" [J. Differential Geom. **35** (1992), no. 1, 85-101], *J. Differential Geom.* **43** (1996), no. 4, 783-788.

[3] J.-C.Birget, S.Margolis, J.Meakin, P.Weil, PSPACE-complete problems for subgroups of free groups and inverse automata, *Theoret. Comput. Sci.* **242** (2000), no. 1-2, 247-281.

[4] O.V.Bogopolski, Finitely generated groups with the M. Hall property, *Algebra and Logic* **31** (1992), no. 3, 141-169.

[5] O.V.Bogopolski, Almost free groups and the M. Hall property, *Algebra and Logic* **33** (1994), no. 1, 1-13.

[6] M. Bridson, D.Wise, Malnormality is undecidable in hyperbolic groups, *Israel J. Math.* **124** (2001), 313-316.

[7] J.J.Cannon, L.A.Dimino, G.Havas, J.M.Watson, Implementation and analysis of the Todd-Coxeter algorithm, *Math.Comp.*, **27** (1973), 463-490.

[8] R.Cremanns, F.Otto, Constructing cannonical presentations for subgroups of context-free groups in polynomial time, Proc. *ISSAC'94*.

[9] M.Dehn, Über unendliche diskontinuerliche Gruppen, *Math. Annn.* **69** (1911), 116-144.

[10] R.Gitik, On quasiconvex subgroups of negatively curved groups, *J. Pure Appl. Algebra* **119** (1997), no.2, 155-169.



Figure 13. The construction of $\Gamma(H_2)$.

[11] R.Gitik, Graphs and separability properties of groups, *J. of Algebra* **188** (1997), no.1, 125-143.

[12] R.Gitik, Doubles of groups and hyperbolic LERF 3-manifolds, *Ann. of Math.(2)* **150** (1999), no.3, 775-806.

[13] R.Gitik, On the combination theorem for negatively curved groups. Corrected reprint of “On the combination theorem for negatively curved groups” [*Internat. J. Algebra Comput.* **6** (1996), no. 6, 751-760]. *Internat. J. Algebra Comput.* **7** (1997), no. 2, 267-276.

[14] R.Gitik, S.W.Margolis, B.Steinberg, On the Kurosh theorem and separability properties, *J. Pure Appl. Algebra* **179** (2003), no. 1-2, 87-97.

[15] M.Gromov, Hyperbolic groups. *Essays in group theory*, 75-263, Math. Sci. Res. Inst. Publ., 8, *Springer, New York*, 1987.

[16] Z. Grunschlag, Algorithms in geometric group theory, PhD thesis, University of California at Berkeley, 1999.

[17] D.F.Holt, Decision problems in finitely presented groups. *Computational methods for representations of groups and algebras (Essen, 1997)*, 259-265, Progr. Math., 173, *Birkhauser, Basel*, 1999.

[18] D.F.Holt, D.Hurt, Computing automatic coset systems and subgroup presentations, *J. Symbolic Computation* **27** (1999), no.1, 1-19.

[19] I.Kapovich, A.Myasnikov, Stallings foldings and subgroups of free groups, *J. Algebra*, **248** (2002), no.2, 608–668

[20] I.Kapovich, A.Myasnikov, P.E. Schupp, V.Shpilrain, Generic-case complexity, decision problems in group theory, and random walks, *J. Algebra* **264** (2003), no. 2, 665-694.

[21] I.Kapovich, A.Myasnikov, P.E. Schupp, V.Shpilrain, Average-case complexity and decision problems in group theory, *Adv. Math* **190** (2005), no.2, 343-359.

[22] I.Kapovich, R.Weidman, A.Miasnikov, Foldings, graphs of groups and the membership problem. *Internat. J. Algebra Comput.* **15** (2005), no. 1, 95-128.

[23] O.Kharlamovich, A.Myasnikov, Hyperbolic groups and free constructions, *Trans. Amer. Math. Soc.* **350** (1998), no.2, 571-613.

[24] O.Kharlamovich, A.Myasnikov, V.Remeslennikov, D.Serbin, Subgroups of fully residually free groups: algorithmic problems, *Contemporary Math*.

[25] N.Kuhn, K.Madlener, F.Otto, Computing presentations for subgroups of polycyclic groups and of context-free groups. *Appl. Algebra in Engrg, Comm. and Comput.*, **5** (1994), no.5, 287-316.

[26] M.Lohrey, G. Senizergues, Rational subsets in HNN-extentions and amalgamated products, in preparation.

[27] R.C.Lyndon and P.E.Schupp, Combinatorial group theory. *Springer-Verlag, Berlin-New York*, 1977.

[28] W.Magnus, A.Karas, D.Solitar, Combinatorial group theory. Presentations of groups in terms of generators and relations. Second revised edition. *Dover Publications, Inc., New York*, 1976.

[29] S.W.Margolis and J.C.Meakin, Free inverse monoids and graph immersions, *Internat. J. Algebra Comput.* **3** (1993), 79-99.

[30] S.W.Margolis, M.Sapir, P.Weil, Closed subgroups in pro-**V** topologies and the extension problem for inverse automata, *Int. J. Algebra Comput.* **11** (2001), no.4, 405-445.

[31] L.Markus-Epstein, Algorithmic Problems in Subgroups of Some Finitely Presented Groups, Phd thesis, Bar Ilan University, 2005.

[32] L.Markus-Epstein, Stallings Foldings and Subgroups of Amalgams of Finite Groups, arXiv.org: math.GR/0705.0754, to appear in *Internat. J. Algebra Comput* (2007).

[33] L.Markus-Epstein, Algorithmic Problems in Amalgams of Finite Groups, arXiv.org: math.GR/0705.0759.

[34] L.Markus-Epstein, Reading Off Kurosh Decomposition, arXiv.org: math.GR/0706.0101 (2007).

[35] J.McCammond, D.Wise, Coherence, local quasiconvexity and the perimeter of 2-complexes, to appear in *Geom. funct. anal.*.

[36] A.Miasnikov, E.Ventura, P.Weil, Algebraic extensions in free groups, arXiv.org: math.GR/0610880 (2006).

[37] C.F. Miller III, On group-theoretic decision problems and their classification. Annals of Mathematics Studies, No. 68. *Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo*, 1971.

[38] C. F. Miller III, Decision problems for groups—survey and reflections. *Algorithms and classification in combinatorial group theory (Berkeley, CA, 1989)*, 1-59, Math. Sci. Res. Inst. Publ., 23, *Springer, New York*, 1992.

[39] A.Minasyan, Separable subsets of GFERF negatively curved groups, *J. of Algebra* **304** (2006), no. 2, 1090-1100.

[40] O.Payne, S.Rees, Computing Subgroup Presentation, Using Arguments of McCammond and Wise, *J. of Algebra*, **300** (2006) (Leedham-Green birthday volume), 109-133.

[41] A.Roig, E.Ventura, P.Weil, On the complexity of the Whitehead minimization problem, arXiv.org: math.GR/0608779 (2006).

[42] P.E.Schupp, Coxeter groups, 2-completion, perimeter reduction and subgroup separability, *Geom. Dedicata* **96** (2003), 179-198.

[43] J.-P.Serre, Trees. Translated from the French by John Stillwell. *Springer-Verlag, Berlin-New York*, 1980.

[44] H.Short, Quasiconvexity and a Theorem of Howson, *Group theory from a geometric point of view*, Proc. ICTP. Trieste, World Scientific, Singapore, 1991.

[45] C.C.Sim, Computation with finitely presented groups. Encyclopedia of Mathematics and its Applications, 48. *Cambridge University Press, Cambridge*, 1994.

[46] J.Stallings, Topology of graphs, *Invent. Math.* **71** (1983), no.3, 551-565.

[47] J Stillwell, Classical topology and combinatorial group theory. *Springer-Verlag, Berlin-New York*, 1980.

[48] N.Tuikan, A fast algorithm for Stallings' folding process, *Internat. J. Algebra Comput.* **16** (2006), no. 6, 1031-1045.

[49] E.Ventura, On fixed subgroups of maximal rank, *Comm. Algebra*, **25** (1997), 3361-3375.

[50] D.Wise, The residual finiteness of negatively curved polygons of finite groups, *Inventiones Mathematicae*, **149**, no.3, 453-685.

Department of Mathematics, Technion, Haifa 32000, Israel
E-mail address: `epstin@math.biu.ac.il`